

VELOCITY FIELD STATISTICS IN STAR-FORMING REGIONS. I. CENTROID VELOCITY OBSERVATIONS

MARK S. MIESCH^a

^aNow at DAMTP, University of Cambridge, Silver Street, Cambridge, CB3 9EW, UK ;
 M.S.Miesch@damtp.cam.ac.uk

Laboratory for Astronomy and Solar Physics
 NASA GSFC, Code 682
 Greenbelt, MD
 miesch@ktaadn.gsfc.nasa.gov
 JOHN SCALO,

Department of Astronomy
 University of Texas at Austin Austin, TX 78712
 parrot@astro.as.utexas.edu

AND

JOHN BALLY
 Center for Astrophysics and Space Astronomy
 University of Colorado, Boulder, Colorado 80309-0389
 bally@casa.colorado.edu

Draft date: February 1, 2008

ABSTRACT

The probability density functions (pdfs) of molecular line centroid velocity fluctuations, and of line centroid velocity fluctuation differences at different spatial lags, are estimated for several nearby molecular clouds with active internal star formation. The data consist of over 75,000 ¹³CO line profiles divided among twelve spatially and/or kinematically distinct regions. These regions range in size from less than to 1 more than 40pc and are all substantially supersonic, with centroid fluctuation Mach numbers ranging from about 1.5 to 7. The centroid pdfs are constructed using three different types of estimators. Although three regions (all in Mon R2) exhibit nearly Gaussian centroid pdfs, the other regions show strong evidence for non-Gaussian pdfs, often nearly exponential, with possible evidence for power law contributions in the far tails. Evidence for nearly exponential centroid pdfs in the neutral HI component of the ISM is also presented, based on older published data for optical absorption lines and HI emission and absorption lines. These strongly non-Gaussian pdfs disagree with the nearly-Gaussian behaviour found for incompressible turbulence (except possibly shear flow turbulence) and simulations of decaying mildly supersonic turbulence. Spatial images of the largest-magnitude centroid velocity differences for the star-forming regions appear less filamentary than predicted by decay simulations dominated by vortical interactions. No evidence for the scaling of difference pdf kurtosis with Reynolds number, as found in incompressible turbulence experiments and simulations, is found. We conclude that turbulence in both star-forming molecular clouds and diffuse HI regions involves physical processes which are not adequately captured by incompressible turbulence or by mildly supersonic decay simulations. The variation with lag of the variance and kurtosis of the difference pdfs is presented as a constraint on future simulations, and we evaluate and discuss the implications of the large scale and Taylor scale Reynolds numbers for the regions studied here.

Subject headings: ISM: clouds, ISM: molecules, ISM: kinematics and dynamics, stars: formation, turbulence

1. INTRODUCTION

Although a great deal of effort has been devoted to quantitatively describing the complex column density spatial structure of star-forming regions (for recent approaches see Falgarone & Phillips 1990, 1991; Gill & Henriksen 1990; Langer, Wilson & Anderson 1993; Adams & Wiseman 1994; Wiseman & Adams 1994; Zimmermann & Stutzki 1993; Houlahan & Scalo 1992; Scalo 1990; Chappell & Scalo 1998; Stutzki et al. 1998), comparatively little attention has been paid to characterising the radial velocity dimension of the data. Exceptions are studies of possible velocity dispersion-size scaling relations (see Falgarone, Puget, & Péroult 1992, Barranco & Goodman 1998, Goodman et al. 1998, and references given there), estima-

tion of the velocity correlation function and related 2-point statistics (see Hobson 1992; Kitamura et al. 1993; Miesch & Bally 1994 and references to earlier work given there) and searches for evidence of rotation (e.g. Goodman et al. 1993). Since one expects a signature of the dynamical and physical processes to appear in the velocity field, and because the velocity field is strongly coupled to, and may in a sense control, the density field and even the star formation rate and the IMF (see models by, e.g. Fleck 1983, Silk 1995, Scalo 1998) it is important to develop additional diagnostics to investigate it. As a step toward a better understanding of molecular cloud velocity structure, Falgarone and coworkers (Falgarone 1989, Falgarone & Phillips 1990, 1991, Falgarone et al. 1994; see below for

discussion) have explicitly tried to relate radial velocity information to dynamical processes through the comparison of observed line profiles with frequency distributions, or probability distribution functions, found in experimental and simulation studies of turbulence. In an earlier short report we (Miesch & Scalo 1995) extended that program to the frequency distribution of line centroid velocities. In the present paper we give a more detailed description and discussion of both the observational data and the centroid probability density functions (which we hereafter refer to as centroid pdfs), and further extend the study to the pdfs of centroid velocity differences. Pdfs of line widths and line skewnesses are examined in a separate paper (Miesch, Scalo & Bally 1999, hereafter Paper II).

The primary goal of this work is to provide useful quantitative observational constraints on ideas, models, and simulations of interstellar turbulence and its relation to star formation—constraints which must be accounted for by any theoretical approach that purports to provide a physical explanatory understanding of the phenomenon. A recurring theme in the present paper is the degree to which interstellar turbulence resembles incompressible turbulence, a field with a very large experimental and theoretical literature. However, we think it is important to recognise from the outset that even if interstellar turbulence *does* turn out to resemble incompressible turbulence in some respects, this would not imply that interstellar turbulence is “understood.” Incompressible turbulence remains an essentially unsolved problem. As many authors have pointed out, there has been (arguably) little tangible progress in the field of incompressible turbulence since Kolmogorov’s seminal paper (Kolmogorov 1941, K41), and most of it consists of elucidating the ways in which K41 was incorrect (e.g. the variation of the scaling exponents of different-order structure functions), and a proliferation of contrasting phenomenological theoretical approaches, mostly imported from other branches of physics (e.g. statistical mechanical and field theoretical approaches; see She 1997 and L’vov & Procaccia 1997). If interstellar turbulence resembles incompressible turbulence, this result may help exclude some physical processes as being dominant, and may implicate the nonlinear advection operator in the momentum equation (which controls incompressible turbulence) as the dominant physical process. However, the physics of this operator in the presence of a large number of degrees of freedom (i.e. large Reynolds number) is still obscure, and one must not confuse the existence of a huge literature with understanding. Furthermore, compressible turbulence does not entail any quadratic invariants (e.g. quantities conserved by the advection operator which are quadratic in the velocity, like kinetic energy), a property which is central to all models of incompressible turbulence. This should from the outset lead one to expect that interstellar supersonic turbulence is different in fundamental ways from incompressible turbulence. From this point of view, the present work is aimed at uncovering observationally these differences.

In §2 we present a discussion of the advantages and disadvantages of the use of centroid pdfs instead of line profiles, and a summary of previous work using this approach (and the velocity difference pdf) for incompressible turbulence, molecular interstellar regions, and extragalactic structure. We also present centroid velocity pdfs

culled from early surveys of optical absorption lines and HI emission and absorption lines in order to show that good evidence already exists for roughly exponential centroid velocity pdfs in the lower-density atomic HI component of the ISM. The data employed in the present study of molecular star-forming regions are presented and discussed in §3, and an overview of the dynamics, star formation activity, and physical environment in each region is provided in Appendix A. In §4 the statistical results are presented, including centroid velocity images and pdfs, as well as velocity difference images, pdfs and pdf moments. Several parametric and non-parametric pdf estimators are used and compared. For the velocity differences, the variation of the pdfs with lag is emphasised and quantified and the possible relation of the non-Gaussian tails to filamentary structures is investigated. The centroid velocity and velocity difference maps are also used to derive effective Reynolds numbers and Taylor scales for the regions considered. The results are summarised in §5. This is the first in a series of papers on the statistical analysis of velocity fields in star-forming regions. In Paper II (Miesch, Scalo & Bally 1999), we use the same data sets presented here to investigate spectral linewidth and line skewness variations, which provide yet another important diagnostic of molecular cloud velocity fields available from densely-sampled emission line observations. In the final paper of the series (Scalo, Chappell, & Miesch 1999, hereafter Paper III), we provide theoretical interpretations of the observed centroid velocity, velocity difference, and linewidth pdfs and a comparison to analytic and numerical models.

2. VELOCITY PDFS AND THEIR ESTIMATION

2.1. General Background

The one-point probability density function contains information that is qualitatively different from correlation functions and related two-point statistics, which are *moments* of some probability density. For example, the autocorrelation function or structure function, while containing spatial information, basically only involve the *variance* of the two-point probability density of velocities as a function of scale. The two-point probability density function itself could be computed, but is difficult to visualise, since, for a given lag scale, it is a two-dimensional surface, and its full representation would involve such surfaces at different spatial lags. More information about the velocity two-point pdf would require high-order moments, which cannot be constructed for ISM data because of noise. In contrast, the one-point velocity pdf, although it contains no spatial information, is easily displayed (it is basically just a one-dimensional histogram), giving access to the probability structure of the velocity field. Similarly, correlations between linewidth and other physical parameters only involve the variance of the one-point pdf, and thus averages over the information in the one-point pdf itself. For these reasons, the one-point velocity pdf offers a qualitatively different view of the velocity field.

The relative independence of the velocity pdf and a second order moment of the two-point pdf, like the power spectrum or correlation function, can be seen in the work of Dubinski, Narayan, & Phillips (1995), who generated simulated line profiles from random velocity fields with a prescribed power spectrum. Although they emphasised

the ability of a Kolmogorov energy spectrum ($k^{-5/3}$) to yield non-Gaussian line-profiles, their results show that other forms of the energy spectrum would yield similar results. In particular, a steeper k^{-2} spectrum, which might be expected for a field of discontinuities or shocks, also gave non-Gaussian profiles of similar form. Thus the 2-point velocity correlation function (a moment) is probably only weakly coupled to the 1-point velocity pdf.

Recent work in several areas suggests that the one-point probability distribution function of dynamical variables like velocity is a useful tool that may be sensitive to dynamical processes. These studies include large scale structure of galaxy velocities (Bernardeau 1994; Kofman et al. 1994; Catelan & Moscardini 1994a,b), incompressible terrestrial turbulence (see below), distinguishing nonlinear chaotic processes from stochastic processes (Wright & Schult 1993), and characterisation of samples of musical volume fluctuations (Scalo & Chappell 1998). In particular, studies of incompressible turbulence have shown that the higher moments (skewness, kurtosis,...) of the pdf can be used to constrain physical models for turbulent intermittency. Although non-zero skewness must exist at some level in order to provide energy transfer among different scales, the pdf of the velocity field itself in incompressible turbulence is in general very nearly Gaussian, at least on large enough scales (Batchelor 1953 and Monin & Yaglom 1971 review early work; see more recent experiments and simulations in Anselmet et al. 1984, Figure 1; Kida & Murakami 1989, Figure 6; Jayesh & Warhaft 1991, Figure 1; Chen et al. 1993, Figure 3.). An important exception is the 3-dimensional incompressible simulation of homogeneous shear flows by Pumir (1996), who finds non-Gaussian, nearly exponential, velocity fluctuation pdfs for velocity components perpendicular to the streamwise component. See also Lamballaise, Lesieur, & Matais (1997) for non-Gaussian velocity pdfs in channel flow close to the boundary.

For incompressible turbulence non-Gaussian behaviour is well-established for velocity *differences* at small scales and velocity *derivatives*, and there is strong evidence from experiments and simulations for non-Gaussian behaviour in many other variables (see the papers referred to above and Chen et al. 1989; Castaing, Gagne & Hopfinger 1990; Vincent & Meneguzzi 1991; She et al. 1993). Often the pdf of the velocity difference or derivative field exhibits a near-exponential behaviour at smaller and smaller scales, and much work has gone into understanding this behaviour physically, especially in terms of the stretching properties of the advection operator (see She 1991 for a review). Part of the motivation of the present work is to investigate whether any of these properties occur in the more complex “turbulence” of interstellar clouds, and whether even the velocity fluctuation field itself presents measurable deviations from a Gaussian pdf.

Falgarone & Phillips (1990, 1991) have shown that line profiles constructed from high-sensitivity CO molecular line data (in several transitions) exhibit excess wing emission, relative to a single Gaussian, over a very large range of scales, from 0.02 to 450 pc. For all these line profiles the width of the wings is about 3 times the width of the line core if both are fit by Gaussians, but the fractional intensity of the wing component (fraction of mass at high velocities) varies between about 0.03 and 0.8. Broad wings

were also found in high latitude molecular clouds by Blitz, Magnani, & Wandel (1988). The presence of similar broad wings in regions whose scales are gravitating and non-self-gravitating, and in regions with and without internal massive star formation, suggests that the behaviour is not due to stellar winds or collapse motions, and the variation in wing width in these regions seems to rule out a dilute warm gaseous component, as pointed out by Falgarone & Phillips. Since the line profile, in the optically thin case, is in effect a histogram of radial velocities, the broad wings have been viewed in the context of non-Gaussian pdfs, although there is some confusion concerning whether the line profile should be interpreted as the pdf of average line of sight velocities or of velocity differences; the latter interpretation is adopted by Falgarone & Phillips (1990) in comparisons with laboratory data.

It is not clear that line profiles give a valid representation of the velocity pdf. Every line profile samples a line-of-sight velocity field which in general contains a component whose characteristic scale is a significant fraction of the sample depth. The form of these systematic line-of-sight motions is unknown and may severely limit the correspondence between the line profile and velocity pdf. Such problems can largely be circumvented in analyses of simulations, where it is possible to insure homogeneity on the largest scales (as in Porter, Pouquet, & Woodward 1994, as analysed by Falgarone et al. 1994), but homogeneity is probably not a good assumption in general for interstellar clouds. It is not difficult to show that the addition of a systematic component can significantly alter the estimate of the distribution of the velocity *fluctuations*, which is the function of interest. A cloud in non-uniform rotation about its center, for example, will yield non-Gaussian line profiles along lines of sight displaced from the projection of the rotation axis onto the plane of the sky (provided this projection is nonzero). In particular, these profiles will exhibit apparent excess wing emission due solely to the smearing arising from the variation of the line-of-sight component of the rotational velocity with depth in the cloud, which will thus distort the pdf of velocity fluctuations. In addition, radiative transfer effects can distort emission lines and cause the wings to become relatively more prominent if the cloud is optically thick (although Falgarone & Phillips 1990 argue against this interpretation of the broad wings on the basis of their observed shapes).

An alternative procedure, which we adopt here, is to estimate the pdf of *centroid line velocities* (intensity weighted average velocity along the line of sight) sampled over a densely observed individual star formation region. While a “line profile” is a measure of the radial velocity (or velocity difference) pdf sampled along the line of sight, either in a single beam or averaged over many beams, the “centroid pdf” is the pdf of the mean velocity of line profiles taken over a large spatial sample of positions in the plane of the sky. The two functions differ in the direction along which the sampling for the pdf is taken, and in the quantity sampled.

The advantages of the centroid pdf approach include the much lower sensitivity required for each of the individual line profiles and the weaker dependence of the results on large scale systematic motions which, although still a concern, will tend to be mitigated by the line-of-sight averaging and by space filtering of the velocity maps (see §3.1).

For example, the centroid velocities of the rotating cloud discussed above will vary in an obvious way with position, and the effects of rotation can therefore be removed by applying an appropriate filter. Such a procedure is not possible with the individual profiles unless the rotation curve of the cloud is known. In addition, the presence of a warm “interclump” medium, or of “optical depth broadening”, which would both contribute to the line profiles, will not much affect the pdf of centroid velocities, since the thermal component and the line saturation are symmetric (although the centroid pdf, in the optically thick case, would only sample fluctuations on the leading edge of the cloud). The problem with this approach is that the number of velocities (positions) which must be sampled in order to accurately estimate the tails of the pdf is very large, at least of order 1000. Furthermore, the relationship between the pdf of an average line-of-sight quantity (centroid velocity in this case) and the pdf of the radial velocity distribution in three dimensions is not clear.

It is also possible to use the centroid velocities to construct the pdf of velocity *differences* for regions separated by a given distance, or, “lag”, and examine how this pdf depends on lag. These velocity difference pdfs are commonly used in studies of incompressible turbulence (see references above), and have been used to study models for the cosmological evolution of galaxy velocities (Peebles 1976; Ueda, Itoh, & Sato 1993; Seto & Yokoyama 1998). It is particularly intriguing that the galaxy velocity difference pdfs exhibit exponential forms at small separations, very similar to the ISM pdfs reported here.

A preliminary account of the velocity difference pdfs obtained for the star-forming regions studied here was presented by Miesch & Scalo (1994). Centroid velocity difference pdfs have also been reported for CO lines in the ρ Oph region (Lis, Keene, et al. 1998; Lis, Phillips, et al. 1998) and for HI lines in the Ursa Major cirrus cloud (Miville-Deschenes, Joncas, & Falgarone 1998). Lis et al. (1996, 1998) find that the velocity difference pdfs in simulations of mildly supersonic decaying turbulence exhibit strong non-Gaussian tails which are associated with filamentary structures and regions of large vorticity, and they report some evidence of such behaviour in molecular cloud observations, at least in quiescent regions. Part of the work presented here is aimed at estimating the pdfs and spatial distribution of velocity differences for regions in which vigorous high-mass star formation has taken place (Orion B and Mon R2 regions) although smaller regions containing only lower-mass YSOs are also represented (L1228, L1551, HH83). It should be recognised at the outset that the pdfs of the centroid velocities themselves for some of these regions already exhibit strongly non-Gaussian tails, (Miesch & Scalo 1995; see also §4.1 below), showing that the “turbulence” in star-forming molecular clouds is different from incompressible and mildly supersonic turbulence, which generally exhibit nearly Gaussian velocity pdfs. Apart from Miesch & Scalo (1995), we know of only one other study in which centroid velocity pdfs of molecular line data have been presented, that of Padoan et al. (1997) who plotted histograms of centroid velocities for several areas in the Perseus cloud, another active star-forming region. Although large-scale motions were not filtered out and the pdf tails were not investigated in detail, these data too clearly exhibit non-Gaussian shapes.

2.2. Previous Estimates for the Atomic ISM

Work aimed at determining the pdf of interstellar gas motions dates back to the early 1950s. Several studies used optical absorption line velocities of “clouds” along the line of sight to OB stars and velocities of HI 21cm emission and absorption lines to estimate the peculiar velocity distribution, after correction for solar motion and differential galactic rotation. These studies refer to fairly local gas, with distances less than about 500-1000 pc. However with the advent and subsequent prominence of molecular line observations, these studies were never repeated and were in effect forgotten.

The results of these earlier studies are presented in Fig.1. In some cases the original histograms were only given in graphical form, and these plots were converted to digitised images using a scanner, and then measured on a computer terminal using standard image manipulation software. We estimate that the uncertainties in reproduction due to this procedure are at the 10 percent level. Since these studies contain a small number of velocities compared to the present work, and because they are only meant to be illustrative of the results already in the literature long ago, the negative velocity portion of the pdf was reflected about $v=0$.

Blaauw (1952) was apparently the first to estimate the form of the velocity pdf, using velocities of CaK lines from Adams’ (1949) catalog. For 150 components along lines of sight to 120 stars with $d < 500$ pc and 80 components toward 43 stars with $d > 500$ pc, and excluding lines associated with the Orion region, Blaauw showed that in both cases there were too many components with high velocity to be fit by a single Gaussian, and that a single exponential could better fit the data. The probable existence of blends due to the relatively poor velocity resolution of the Adams survey was a problem, and Blaauw tried to correct his pdfs using an ingenious technique that is notable for its use of humans as an analog random number generator. The resulting pdf for the 150 nearby components (Blaauw’s Table 5a) is shown in Fig. 1 (filled circles) in log-linear form. Notice that in this form a Gaussian would be a parabola and an exponential would be a straight line. Takakubo (1967) also examined the K line velocities of Adams and concluded that an exponential pdf gave a better fit than a Gaussian. Huang (1950) and Kaplan (1954) preferred $1/v$ fits to the optical line pdf, based again on Adams’ catalogue. (See Fig. 1 in Kaplan 1966). Siluk & Silk (1974) examined the pdf of the high-velocity ($v > 20 \text{ km s}^{-1}$) tail of the Adams optical line sample and fit the pdf with a power law of index around -3 . Munch (1957) presented new observations of optical interstellar lines, at poorer resolution, toward 132 stars. He used the “doublet ratio method” (CaH/K or NaD2/D1) to compare curves of growth with calculations based on Gaussian and exponential velocity pdfs. The Gaussian fit required a physically unrealistic increase of velocity dispersion with distance, while the exponential provided a fit with a single mean speed, confirming Blaauw’s conclusion.

Blending remained a problem with these interpretations. Hobbs used interferometric scans of NaD (Hobbs 1969a,b) and KI (Hobbs 1974) in an attempt to fit individual profiles, and showed that most of Adams’ lines were probably multiple. Hobbs found that the majority of individual line

profiles could be fit by Gaussians, while a smaller fraction favoured an exponential, and many lines were not well-fit by either distribution; however the velocity pdf was not examined. Falgarone & Lequeux (1973) used Hobbs’ NaD data to estimate the cloud peculiar velocity dispersion, but the number of components was too small for an estimate of the pdf. Later work on optical interstellar lines turned almost exclusively toward studies of interstellar depletion patterns, and the question of the velocity pdf was never re-examined.

FIG. 1.— Histogram representation of the radial velocity probability density function (pdf) for distinct velocity components identified in optical absorption line surveys along the line of sight to OB stars and HI emission and absorption line surveys. The data were either taken from tables in the papers indicated, or measured from digitised scans of graphs presented in the papers. The negative-velocity portions of the pdfs have been reflected about $v = 0$. In this log-linear plot a Gaussian pdf would appear as a parabola while an exponential pdf would be a straight line. Additional studies not shown in this figure are discussed in the text.

Meanwhile, HI 21cm survey data was accumulating. Takakubo (1967) performed Gaussian decomposition and subtraction of solar motion and galactic rotation, resulting in peculiar velocities for 544 HI emission lines whose velocity pdf was presented for 3 intervals of linewidth and different galactic latitude intervals. He judiciously refrained from any conclusions concerning the form of the pdf because of the dependence of the derived velocity dispersions on galactic latitude. (This was perhaps the first paper to suggest consistency with the Kolmogorov velocity spectrum for incompressible turbulence, on the basis of the power law scaling of velocity dispersion with the sine of the latitude corresponding to a scaling of the structure function with path length.) The resulting histogram for all 544 velocities is nevertheless shown in Fig. 1 (reflected about $v=0$) as filled inverted triangles. Mast & Goldstein (1970) presented a similar analysis for 268 peculiar radial velocities of HI 21 cm emission clouds that were well-resolved in velocity. Their Figure 4 clearly shows that an exponential fits better than a Gaussian: a single Gaussian can fit the wings or the core but not both. Their observed pdf is shown in Fig. 1 (open circles).

Crovisier (1978) studied the velocities of HI clouds identified as components of Gaussian decomposition of absorption profiles observed towards extragalactic sources in the Nançay survey. About 300 such clouds were selected on the basis of latitude, so that they might represent relatively local gas. The histogram of residual velocities, after corrections for solar motion and differential galactic rotation, is given in Crovisier’s Fig. 1. The interpretation of the resulting histogram of residual velocities is problematic because of the possibility that the intermediate velocity features are spurious. The histogram for all the components is displayed in Fig. 1 (filled squares). Dickey, Salpeter, and Terzian (1978) presented the residual velocity distribution of clouds from the high-sensitivity and high-resolution Arecibo 21 cm emission/absorption survey, but the number of velocities is too small to estimate the

functional form of the pdf. Further discussion of the high-velocity tail of the HI distribution and implications for turbulent energy requirements can be found in Kulkarni & Fich (1985) and Lockman & Gehman (1991).

All of the above studies included lines of sight that cover a significant fraction of the sky and a range in distance, rather than focusing on individual cloud complexes. An exception is the detailed HI emission mapping of two regions by Verschuur (1974) using the NRAO 300 ft. antenna. Identifying “clouds” as distinct entities in spatial-radial velocity space, Verschuur presented histograms of velocities for about 150 clouds in his region A and about 50 in region B. The region B histograms cannot be used to determine the intrinsic pdf of velocity fluctuations without careful filtering (which we have not attempted) because of ordered motion along the filamentary structures in the region. The histogram for region A, reflected about $v=0$, is shown as open squares in Fig. 1.

It should be emphasised that the rotation curve of the Galaxy and other large-scale trends have been subtracted out in all of the studies mentioned above. Furthermore, the results are likely not influenced by the Local Bubble, since the Verschuur sample was for a single region at a well-defined position and velocity, the optical-line clouds were detected toward OB stars which are on average about a kpc away, and the HI emission and absorption studies likewise sample much larger distances (the Local Bubble only extends about 100 pc from the sun). It is possible that the optical line studies are influenced by gas motions in local HII regions around the OB stars, so some caution should be taken in their interpretation. However, it is not clear that the clouds sampled in such studies lie in general at the same distance as the stars themselves (many studies have suggested otherwise), and the pdfs derived from the optical line data are similar to those derived from HI data, which are certainly not biased toward hot stars.

Overall, it can be seen from Fig. 1 that evidence for non-Gaussian, and probably exponential, centroid velocity pdfs in the relatively low-density atomic HI component of the ISM has existed for some time. Many of these earlier works had already concluded that the velocity pdf is better fit by an exponential (or $1/v$) than a Gaussian. It is surprising, and unfortunate, that all these early studies have been in effect “forgotten” in more current discussions of interstellar turbulence, since they suggest that at least some forms of interstellar turbulence differ significantly from incompressible turbulence, for which the velocity pdf is invariably nearly-Gaussian. The implication is that the physical processes involved are substantially different than for incompressible turbulence. This result for relatively low-density HI gas has recently received considerable support from the detailed aperture synthesis HI study of the entire LMC by Kim et al. (1998). For scales above the resolution limit of 15pc, the peculiar (fluctuation) velocity pdf of the gas (for motions out of the plane of the LMC) is extremely well-fit by an exponential, and a Gaussian seems to be certainly excluded.

We note that most of the HI studies reviewed in this section concern clouds which are probably not actively forming stars. So there is good evidence that, even in some “quiescent” regions, interstellar gas motions exhibit properties inconsistent with experiments and simulations of incompressible turbulence. On the other hand, Falgar-

one et al. (1994) have found good agreement between the shapes (as quantified by moments of order 2 and 4) and point-to-point shape variations of observed spectral line profiles in quiescent molecular clouds and synthesised profiles from the “nearly incompressible” phase in simulations of mildly supersonic, decaying turbulence by Porter, Pouquet, & Woodward (1994). Apparently, either the physical processes occurring in at least some quiescent molecular clouds and HI clouds are substantially different, or the probability density function of centroid velocities contains different information than the second and fourth-order moments of individual line profiles. The former possibility is likely, since the HI linewidths reported in the papers discussed above are significantly supersonic in most cases.

We also point out that shapes and shape variations in optically thin line profiles similar those reported by Falgarone et al. (1994) for mildly compressible turbulence simulations have also been claimed for completely non-dynamical incompressible velocity fields (Dubinski, Narayan, & Phillips 1995), highly supersonic simulations of cloud collisions (Keto & Lattanzio 1989), and highly supersonic stellar-driven self-gravitating turbulence fields by Vazquez-Semadeni et al. (Vazquez-Semadeni, private communication). For this reason we are not convinced that the shapes and shape variations of individual line profiles alone can be used to discriminate between models, even for relatively quiescent clouds. Additional diagnostics of the velocity field are therefore needed.

This section has been a review of centroid velocity pdfs observed in the diffuse atomic phase of the interstellar medium. In the remainder of this paper, we extend these studies to the higher-density molecular gas in active star-forming regions and also consider the pdf of velocity differences. Ultimately our goal is to use these statistics of the velocity field to help understand the physical processes responsible for interstellar turbulence.

3. OBSERVATIONS

The observations used in this paper are the same as those used in the previous statistical analyses of Miesch & Bally (1994, hereafter MB) and Miesch & Scalo (1995, hereafter MS), and include large-scale mappings of the giant molecular clouds in the northern part of the Orion region (Orion B) and in the Monoceros region, associated with the Mon R2 infrared cluster. Also included are observations of the smaller-scale clouds which appear in the Lynds catalog as numbers 1228 and 1551 (Lynds 1962), as well as observations of the molecular gas surrounding the Herbig-Haro object HH83. In what follows, as in MS and in MB, we will refer to these five separate mappings as Orion B, Mon R2, L1228, L1551, and HH83.

The Orion B, Mon R2, L1228, and L1551 observations are emission-line measurements of the 110 GHz, $J = 1 \rightarrow 0$ transition in ^{13}CO . They were obtained with the AT&T Bell Laboratories 7 m offset Cassegrain antenna located in Holmdel, New Jersey, between 1985 December and 1991 June. This antenna has a very clean Gaussian beam with a FWHM of $100''$ between 98 and 115 GHz and the effective channel bandwidth (resolution) is 100 kHz (0.27 km s^{-1}) for the Orion B, Mon R2, and L1228 data sets and

50 kHz (0.14 km s^{-1}) for the L1551 data set. Typical signal-to-noise ratios are between 5 and 9. Further details on the observations and measurement techniques can be found in Bally, Langer, Stark, & Wilson (1987), Bally, Stark, Wilson, & Henkel (1987), Bally, Stark, Wilson, & Langer (1989), Pound, Bania, & Wilson (1990), Bally et al. (1991), Pound & Bally 1991 (1991), Bally et al. (1995), and MB.

The observations of the HH83 molecular cloud were obtained with the IRAM-30m telescope on Pico Veleta, Spain in April and December 1989, and have a higher spatial resolution (half power beamwidth = $13''$) and smaller field of view. The mapped region lies just west of the Orion A molecular cloud (L1641) and is associated with the Herbig-Haro object HH83. The measurements are of the ^{13}CO $J = 2 \rightarrow 1$ emission line at 220 GHz, with an effective frequency resolution of 100 kHz (0.13 km s^{-1}). A typical signal-to-noise ratio for these observations is about 20. More information on the data acquisition, as well as the structure, environment, and physical conditions in the molecular cloud can be found in Bally, Castets, & Duvert (1994).

We recognise that ^{13}CO is an imperfect tracer of the gas distribution and velocity field in dense molecular clouds. The primary problem is the likelihood that the ^{13}CO transitions are optically thick along lines of sight with the largest total column density. Studies of the correlation of ^{13}CO line strength with visual and infrared extinction and with rarer CO species such as C^{18}O (Frerking, Langer, & Wilson 1982; Frerking et al. 1989; Lada et al. 1994) suggest that ^{13}CO saturates at visual extinctions greater than about 5 mag, although these papers only study a small number of individual clouds. Furthermore, the comparison of C^{18}O emission with infrared extinction by Alves, Lada, & Lada (1998) suggests that for $A_v \gtrsim 10$ mag, C^{18}O underestimates the total optical depth, possibly due to depletion of CO onto grain surfaces. These results imply that the present observations are biased in the sense that they do not penetrate the highest-column density substructure. However, the fractional area covered by column densities greater than $A_v \approx 5$ is small, about five or six percent in Orion B, one or two percent in Mon R2, and effectively zero in L1228, L1551, and HH83 (assuming the following conversion factors: $\text{I}(^{13}\text{CO}) \rightarrow \text{N}(^{13}\text{CO}) = 5.9 \times 10^{14} \text{ cm}^{-2} \text{ K}^{-1} \text{ km}^{-1} \text{ s}$, $\text{N}(^{13}\text{CO}) \rightarrow \text{N}(\text{H}_2) = 7 \times 10^5$, and $\text{N}(\text{H}_2) \rightarrow A_v = 1.06 \times 10^{-21} \text{ mag cm}^2$). So, we are probably probing *most* of the molecular gas in the surveyed regions. Furthermore, it would be extremely time-consuming to collect, say, C^{18}O spectra for the large number of lines of sight (more than 75,000) examined here, because of the required sensitivity. So, in terms of spatial coverage and statistical characterisation of the velocity field, ^{13}CO is probably the best probe available. Nevertheless, the fact that this tracer probably does not probe the highest-column density substructure should be kept in mind in interpreting the statistical results of §4 below.

The observations are presented in Figures 2 and 3. The coordinates corresponding to the origin (0,0) of each map are listed in Table 1 (see also MB, Table 1). The centroid velocity¹ along each line of sight is indicated by the color

¹The centroid velocity in each line of sight is defined as $\Sigma T_i v_i / \Sigma T_i$, where T_i and v_i are the brightness temperature and velocity corresponding to the i^{th} spectrometer channel, and the summations span some selected velocity range.

TABLE 1
OBSERVED CLOUDS

Region	R.A. (1950)	Decl. (1950)	Distance ^a (pc)	Size (pc)	Mass ^a ($M_{\odot}$)	Mach Number ^b
Orion B	05 ^h 41 ^m 08 ^s .1	−01° 05′ 00″	415 [1]	20×20 ^c ; 45×20	8×10 ⁴ [6]	1.9–2.9 ^c ; 4.0–7.1
Mon R2	06 05 22.0	−06 22 25	830 [2]	45×40	9×10 ⁴ [6]	2.6–3.9
L1228	20 58 00.0	+77 23 00	300 [3]	2.6×2.6	200–900 [7]	2.6
L1551	04 28 40.0	+18 01 42	140 [4]	2.1×1.5	80 [8]	1.7
HH83	05 31 06.3	−06 31 45	480 [5]	0.9×0.5	8–15 [9]	1.5

^aReferences are as follows: [1] Anthony-Twarog 1982, [2] Racine 1968, [3] Grenier et al. 1989, [4] Elias 1978, [5] Genzel et al. 1981, [6] Maddalena et al. 1986, [7] Bally et al. 1995, [8] Snell 1981, [9] Bally, Castets, & Duvert 1994

^bThese Mach numbers are based on the centroid velocity fluctuation amplitudes listed in Table 5 (σ_c), and an assumed kinetic temperature of 20K.

^cThe first values listed correspond to regions 1a, 1b, and 1c, while the second values represent the remainder: regions 2, 3, and 4 (see below).

TABLE 2
REGIONS ANALYSED

Region	Number of Spectra	Grid Spacing (arcmin)	Interpolation Radius (arcmin)	Number of Points	LSR Velocity Integration Limits (km s ^{−1})
Orion B (1a)	943	0.5	1.1	822	[9.35,11.75]
(1b)	1604	0.5	1.1	2094	[7.5,10.5]
(1c)	5817	0.5	1.1	4801	[8.5,12.5]
(2)	25510	0.5	1.1	18420	[5.0,14.5], [9.2,11.9], [7.0,12.1]
(3) ^a	23936	0.5	1.1	10433	[3.0,9.2], [1.0,7.5]
(4)	25311	0.5	1.1	21876	[7.5,12.7]
Mon R2 (1)	3260	0.5	1.1	8630	[10.6,13.4]
(2)	2813	0.5	1.1	7592	[7.7,12.5]
(3)	4570	0.5	1.1	12208	[11.2, 14.4]
L1228	1922	0.5	1.1	3532	[5.5,8.1]
L1551	2327	0.5	1.1	4726	[−9,−6]
HH83	1142	0.05	0.25	6235	[5.0,7.5]

^aAs discussed in the text, region 3 in Orion B overlaps spatially with regions 2 and 4 and therefore includes most of the same spectra, but integrated over a different velocity range.

image, with blue and red denoting movement respectively toward and away from the observer with respect to the mean LSR velocity of the cloud. The contour plot overlays indicate the integrated intensity in the ¹³CO transition considered (see above).

The number of independent spectra included in these observations totals more than 75,000 (see Table 2). The spectra for each region were extracted, reduced, and interpolated onto a regular spatial grid as described by MB. The grid spacing and interpolation radius used for each

set of observations is listed in Table 2. The velocity integration ranges for each region, also listed in Table 2, were chosen to be large enough to span most of the emission, but still small enough to mitigate the influence of instrumental noise on the centroid fluctuations. In the case of Orion B and Mon R2, some of the integration ranges were further restricted in order to isolate particular cloud components. See MB for further details.

After interpolating the spectra onto regular spatial grids, threshold integrated intensity levels were introduced

FIG. 2.— Shown are the Orion B $^{13}\text{CO } J = 1 \rightarrow 0$ observations used in the analysis to follow. Contours represent integrated intensity and images represent centroid velocity. Red and blue in the centroid velocity images correspond respectively to receding and approaching motions relative to the mean LSR velocity of the cloud (see the color table in Fig. 3). The data set has been subdivided into 6 distinct regions as described in the text. The orientation of the coordinate system with respect to the north and west directions on the plane of the sky is indicated. All coordinates represent the offset in arcminutes from the position of the origin, which is listed in Table 1 and applies to all regions. The dashed blue line indicates the position of region 3, which overlaps regions 2 and 4. These, together with the dotted white lines, also serve to delimit the composite portions of regions 2 and 3 which were obtained using different integration limits as described in the text. The LSR velocity integration limits used for each region are indicated in brackets

FIG. 3.— Similar to Figure 2, but for the $^{13}\text{CO } J = 1 \rightarrow 0$ Mon R2, L1228, and L1551 observations and the $^{13}\text{CO } J = 2 \rightarrow 1$ HH83 observations. The color table used for the centroid velocity images (which applies also to Fig. 2), is displayed on the right. The orientation of the coordinate system is indicated in each panel and all axes represent arcminute offsets with respect to the positions listed in Table 1.

in order to eliminate pixels with low signal-to-noise. Any remaining low-intensity emission located outside the main cloud boundaries and having dramatically different centroid velocity values (likely influenced by noise) was also removed. The number of pixels in each of the resulting centroid velocity maps in Figures 2 and 3 are listed in Table 2. Since the maps were oversampled, the number of points in the centroid and integrated intensity maps is in many cases larger than the associated number of independent spectra.

Several distinct cloud components, or “regions”, distinguished by their spatial position and their mean LSR velocity, emerge when the Orion B and Mon R2 data cubes are analysed closely. In the study presented here, as in MS and in MB, we have divided these data sets accordingly and have considered each of the components separately. Thus, in Figures 2 and 3a we have indicated six regions (1a, 1b, 1c, 2, 3, and 4) in Orion B and 3 regions (1, 2, and 3) in Mon R2. In order to facilitate the isolation of specific cloud components in the Orion B and L1551 observations, we have chosen coordinate systems which are tilted somewhat with respect to the right ascension - declination standard. The orientation of each map is indicated by arrows in Figures 2 and 3.

All the coordinate axes in Figure 2 are with respect to the same origin in order to facilitate the placement of the different regions for the reader. Thus, region 1 (a, b, and c), lies directly above region 2 in the coordinate system chosen (see MB, Fig. 1e), or in other words, just north-east of the main Orion B cloud. It has been displaced and enlarged somewhat in Figure 2 for clarity. Region 3 overlaps regions 2 and 4 in the manner indicated by the dashed blue lines. It appears to be a fairly well-defined separate cloud component which lies at a significantly lower LSR velocity (by $\sim 4 \text{ km s}^{-1}$) than the remaining emission and was isolated by choosing appropriate velocity integration limits (see below and MB). It is likely associated with the doubly-peaked ^{12}CO line profiles in this region

reported by Maddalena et al. (1986). In order to isolate the spatially overlapping regions in Orion B, we further subdivided regions 2 and 3 into portions, each integrated over a somewhat different velocity range, as indicated in Figure 2. These portions were then combined to produce composite maps for regions 2 and 3.

The cloud components used for Mon R2 are shown in the three separate panels of Figure 3a. As in the case of Orion B, they were chosen on the basis of both spatial connectivity and mean LSR velocity using channel maps and spatial-velocity diagrams. The relative orientation of these regions can be discerned by noting the coordinate axes in each panel (which are all with respect to the same origin) and by referring to Figure 1d of MB. The distinction between the three different regions defined here is also apparent in the ^{12}CO channel maps presented by Xie & Goldsmith (1994; their Figures 1 and 2), who interpret the relative blueshift of the southeastern portion of the cloud (our region 1) as evidence for a large-scale, expanding shell (see §A.2 in Appendix A).

An overview of the physical conditions and environment in each of the molecular clouds in our study is given in Appendix A. Briefly, our sample covers a broad range of scales and conditions relevant to the velocity field, with energy input ranging from hot massive stars to low-mass protostellar outflow sources. What the regions have in common is that they all contain strong, active sources of momentum and energy to drive the “turbulence” whose statistics we wish to study.

For reference, Table 1 lists estimates for the distance to each region, and for the total mass and plane-of-the-sky extent of the CO-emitting gas. Also listed are estimates for the Mach number of each region, based on the measured rms centroid velocity fluctuation (see Table 5 below) and on an assumed sound speed of 0.287 km s^{-1} (corresponding to a kinetic temperature of 20K). Note that the range is wide, from 1.5 to 7.1, and that all regions are substantially supersonic, suggesting that compressibility may play an

important role in their dynamics.

4. STATISTICAL ANALYSIS

4.1. Centroid Velocity Fluctuations

4.1.1. Analysis

It is apparent in the centroid velocity images shown in Figures 2 and 3 that many of the regions studied have large-scale velocity gradients across them, which at least in some cases, dominate any statistical quantifiers which may be computed. Since statistical approaches generally lose spatial information by considering only distributions or averages, their utility depends on some degree of uniformity in the data. Any statistical quantifier which is dominated by a small number of large-scale features in the data can give misleading results. In other words, large-scale gradients should be removed from the centroid data in order for our statistical results to be meaningful and comparable to theoretical models, numerical simulations, and laboratory experiments involving turbulent fluids.

For these reasons, we have removed large-scale gradients by first applying a low-pass filter, or smoothing function, to each data set and then subtracting this smoothed map from the original to obtain the centroid velocity fluctuations. The filters were chosen to be as wide as possible, while still eliminating any prominent, extensive, antisymmetric “lobes” in the autocorrelation function of the fluctuation maps. Such features in the autocorrelation function are the characteristic signature of large-scale gradients in the data. The details of the filtering process are described by MB and will not be discussed further here.

All of the analysis presented in this and the following sections (§4.1–§4.3) is based on the centroid velocity fluctuations, which are shown as Grey-scale images in Figures 4 and 5. Although the pdfs of the centroid velocity fluctuations (discussed below) generally appear significantly different from the unfiltered data, they are not sensitive to the precise value of the effective cutoff wavelength or the shape of the filter. Furthermore, since the analysis of velocity differences in §4.2 primarily samples small-scale structure, it is very insensitive to spatial filtering.

Notice that although some of the highest-amplitude fluc-

tuations, which will appear in the far tails of the corresponding pdfs, are found within the main bodies of the mapped regions, others occur near the cloud edges or in “outlier” regions—areas which could not be observed over large continuous spatial extents due to the sensitivity limits of the observations or the influence of overlapping cloud components. For example, this behaviour can be seen in the L1551 and Orion B, region 1b maps in Figures 4c and 5e. Such high-amplitude edge or outlier points introduce a bias in the far tails of the estimated pdfs if they represent emission that is noise-dominated or kinematically distinct from the main body of each region. Although much of this emission was removed from the maps prior to analysis, it is very difficult to identify and remove all of it. The centroid velocity pdf estimates therefore still contain some of this bias. The influence of noise and overlapping cloud components is discussed further in Appendix B.

The pdfs, or probability density functions, corresponding to each of the centroid velocity fluctuation images of Figures 4 and 5 are shown in Figures 6 and 7. Although the classical histogram is a familiar and robust estimator of the pdf, it is generally not the most accurate. Vio et al. (1994) have investigated several alternative pdf estimators in addition to the histogram and have compared their performance on some typical astronomical problems. They conclude that the histogram is decidedly inferior in the applications they considered and recommend the use of more sophisticated pdf estimators. Motivated by this result, we have implemented two of the histogram alternatives described by Vio et al. (1994): the adaptive kernel estimator and the Johnson estimator.

Kernel estimators approximate the pdf at each data value in terms of a sum over all data points, weighing each according to a specified window function, or kernel. They do not require binning of the data as in the classical histogram estimator and are generally less sensitive to noise in the data, producing a smoother, continuous, and often more reliable result. In the adaptive kernel method, the effective width of the kernel increases systematically where the data density is low, thereby improving accuracy in the pdf tails. In our implementation, the form of the kernel and the manner in which its width is allowed to

FIG. 4.— Shown are grey scale images of the centroid velocity fluctuations in Orion B, which were obtained by filtering the raw centroid velocity maps shown in Figure 2 as described in the text. Increasing brightness corresponds to a velocity directed increasingly away from the observer, as indicated by the intensity table in Fig. 5.

FIG. 5.— Similar to Figure 4, but for the Mon R2, L1228, L1551, and HH83 data sets. These maps were obtained by applying spatial filters to the centroid velocity maps shown in Figure 3 (see text). The sense of the grey scale is shown by the intensity bar, which also applies to Figure 4.

vary are as described by Vio et al. (1994). The adaptive kernel estimation of each of the pdfs in Figures 4 and 5 is shown as a dashed line.

An alternative strategy for pdf estimation is based on defining some general class of analytic functions involving several parameters which can be approximated using statistical moments or percentiles derived from the data set. The advantage of such an approach is that it does not require the subjective binning of data or the choice of a kernel function and window width. It can also be less sensitive to noise in the data than either histogram or kernel methods. The disadvantage of parametric methods is that, however general, they impose some functional form to the data which may not be present. Our implementation of the Johnson system follows that described by Vio et al. (1994). Since the centroid velocity is in principle an unbounded variable, we consider only the so-called S_U family of Johnson estimators, which is characterised by the four parameters η , ϵ , λ , and γ , and which is defined by the following functional form:

$$f_J(v) = \frac{\eta}{\sqrt{2\pi}[(v-\epsilon)^2 + \lambda^2]} \times \exp \left\{ -\frac{1}{2} \left[\gamma + \eta \ln \left(\frac{(v-\epsilon)}{\lambda} + \sqrt{\left(\frac{(v-\epsilon)}{\lambda} \right)^2 + 1} \right) \right]^2 \right\} \quad (1)$$

for $-\infty < x < \infty$. The Johnson estimation for each of the pdfs in Figures 4 and 5 is shown as a solid line, and the best-fit parameters are listed in Table 3.

The centroid velocity pdfs presented by MS apply to the same data as the pdfs in Figures 6 and 7, but they were based solely on the histogram pdf estimator (with a slightly different binning than that used here). MS also constructed several composite pdfs, one of which was formed from the six subregions of Orion B, another from the three subregions of Mon R2, and a third composite pdf which was composed of all twelve data sets. For the sake of brevity, we have chosen to omit these composite pdfs from the present study, although it is worth noting that the Orion composite pdf in particular appears very nearly exponential.

Most of the pdfs in Figures 6 and 7 exhibit significantly non-Gaussian shapes, a result which will be discussed in §4.1.2 below. However, before proceeding, we note that a few pdf features probably originate from the superposition of two or more distinct cloud components. For example, the morphology and kinematics of regions 1b and 1c in Orion B (see Figs. 2 and 4) suggest several cloud components, which may be the source of some of the bimodality and fine structure exhibited by the associated pdfs (Figs. 6b and 6c). Although the spatially overlapping cloud components in Orion B regions 2 and 3 were separated out by choosing appropriate velocity integration limits, some contamination remains. Thus, the low velocity ($< -2 \text{ km s}^{-1}$) pdf tail in region 2 (Fig. 6d) and the associated large kurtosis value of 11 (see Table 4 below), are probably influenced by residual emission from region 3, and the abrupt high-velocity ($\sim 2 \text{ km s}^{-1}$) cutoff in the region 3 pdf (Fig. 6e) is likely influenced by the imposed upper integration limit. Also, residual overlap in Mon R2 may account for some of the excess in the high-velocity tail of region 1 and the low-velocity tail of region 2 (Figs. 7a and b).

Another way to quantify the spread and shape of probability density functions is by the sample moments, computed directly from the data set. We consider the first four central moments, defined as follows:

$$\text{mean} = \mu = \frac{1}{N} \sum_{x,y} v(x,y) ; \quad (2)$$

$$\text{standard deviation} = \sigma = \sqrt{\frac{1}{N} \sum_{x,y} [v(x,y) - \mu]^2} ; \quad (3)$$

$$\text{skewness} = \frac{1}{\sigma^3} \frac{1}{N} \sum_{x,y} [v(x,y) - \mu]^3 ; \quad (4)$$

$$\text{kurtosis} = \frac{1}{\sigma^4} \frac{1}{N} \sum_{x,y} [v(x,y) - \mu]^4 . \quad (5)$$

The summation over x, y spans the area of the cloud or in other words, the entire data set. The first two moments quantify the location and spread of the pdf, while the third and fourth are nondimensional quantities which contain information about its shape. In particular, the skewness and kurtosis are measures of the symmetry and

FIG. 6.— Centroid velocity pdfs for Orion B, corresponding to the images shown in Figure 4. Crosses denote the histogram estimate with statistical ($N^{1/2}$) error bars and solid and dashed lines represent the Johnson and adaptive kernel estimators discussed in the text. The corresponding Johnson parameters are listed in Table 3.

FIG. 7.— Similar to Figure 6, but for the Mon R2, L1228, L1551, and HH83 data sets shown in Figure 5. The parameters corresponding to the Johnson estimators (solid lines) are listed in Table 3.

TABLE 3
JOHNSON PARAMETERS FOR THE CENTROID VELOCITY PDFS

Region	η	ϵ	λ	γ
Orion B (1a)	1.55	-1.63×10^{-2}	0.192	-0.114
(1b)	5.20	-0.292	1.01	-0.292
(1c)	1.17	0.218	0.285	0.504
(2)	2.03	0.149	0.534	0.670
(3)	2.08	-0.318	1.13	-0.604
(4)	1.32	-7.75×10^{-2}	0.476	-0.157
Mon R2 (1)	5.37	1.26	1.46	4.31
(2)	4.27	0.121	0.929	0.609
(3)	2.80	-6.70×10^{-2}	0.769	-0.349
L1228	1.64	5.00×10^{-2}	0.175	0.322
L1551	1.11	-5.65×10^{-2}	0.141	-0.201
HH83	1.59	5.29×10^{-4}	0.158	0.255

“flatness” of the pdf respectively. A Gaussian distribution has a kurtosis of 3, and a value larger than 3 implies that the pdf in question has relatively more prominent tails – in other words, high-amplitude events are more numerous than would be expected for a Gaussian random variable. In the turbulence literature this behaviour is usually referred to as “intermittency” (although the same term is often used in reference to the variation of the scaling exponents of the moments of the velocity *difference* distribution). A kurtosis less than three implies the opposite—that pdf tails are less prominent relative to a Gaussian distribution. An exponential distribution ($f(x) \propto \exp(-|x|)$) has a kurtosis equal to 6. Table 4 lists the second, third, and fourth moments for each of the pdfs shown in Figures 6 and 7. Note that the skewness and kurtosis values listed are central sample moments, and are therefore slightly different from the values given by MS, which were computed using the histogram as a pdf estimator.

Many theoretical and empirical studies involving pdfs in the context of incompressible turbulence are concerned with stretched exponential forms, whereby the pdf tails fall off as $f(x) \propto \exp\{-a|x|^\beta\}$ (e.g. Kailasnath, Sreenivasan, & Stolovitzky 1992; Lohse & Grossmann 1993; Yee & Chan 1997; Frisch & Sornette 1997). A Gaussian pdf is characterised by $\beta = 2$, and an exponential pdf by $\beta = 1$. Fractional β values can arise from random multiplicative processes and represent a straightforward and useful way of quantifying the departure of a pdf from Gaussian statistics.

The value of the stretching exponent β determines how quickly the pdf decays, and as a result, how prominent the pdf tails are. For a properly normalised stretched exponential pdf, there is a one-to-one correspondence between the value of β and the pdf kurtosis. Therefore, if the observed velocity centroid pdfs can be approximately described by stretched exponential forms, then the sample kurtosis values listed in Table 4 imply an effective β for each data set. The results, listed in Table 4, yield stretching exponents for Orion B and the smaller clouds (L1228,

L1551, and HH83) which generally range between 1.0 and 1.4. Exceptions include Orion B regions 1a ($\beta = 2.1$) and 2 ($\beta = 0.71$). The Mon R2 regions yield higher β values, between 1.8 and 2. Curve fits to the tails of the Johnson pdf estimators yield similar, although slightly lower results for the stretching exponents. Adopting a Poisson weighing and combining the high and low velocity tails to improve statistics, such curve fits, excluding Mon R2, yield β values ranging from 0.77 to 1.2, with the exception of Orion B, region 1b ($\beta = 1.6$). The results for Mon R2 again indicate somewhat steeper tails, with $1.5 < \beta < 1.7$.

We have also considered the possibility that power law forms, $f(x) \propto x^n$, may fit at least some of the pdf tails better than stretched exponentials. Power law velocity distributions could arise from stellar winds, outflows or superbubbles (e.g. Silk 1995; Oey & Clarke 1998) and would suggest physical processes substantially different than those occurring in isotropic, incompressible turbulence. When the pdf tails of Figures 6 and 7 are plotted on log-log axes rather than log-linear axes (Fig. 8), many of them appear linear in portions, suggesting power laws. This is particularly the case for L1228, HH83 and Orion B, region 4, and possibly also in L1551 and Orion B, regions 1a, 2, and 3. Curve fits to the pdf estimators in these regions show some evidence for power law indices of $\approx -4.5 \pm 1$, although the positive velocity tails in L1551 and Orion B, region 3 seems significantly shallower ($n \approx 2$ or 3). Power law forms are displayed in each of the frames of Fig. 8 for comparison. Although stretched exponentials in most (but not all; c.f. L1228) cases appear to provide somewhat better fits to the observed pdf tails than power laws, the data are generally not of high enough quality to make a reliable distinction.

4.1.2. Discussion

In most cases, there is good agreement between the three different pdf estimators (histogram, adaptive kernel, and Johnson) shown in Figures 6 and 7. Exceptions include the pdf excesses in the tails of Orion B regions 1b and 2,

TABLE 4
MOMENTS AND STRETCHING EXPONENTS FOR THE CENTROID VELOCITY PDFS

Region	Standard Deviation (km s ⁻¹)	Skewness	Kurtosis	β
Orion B (1a)	0.15	-0.018	2.9	2.1
(1b)	0.22	-0.50	4.8	1.2
(1c)	0.36	-0.58	4.2	1.4
(2)	0.34	-1.3	11	0.71
(3)	0.63	0.073	4.0	1.4
(4)	0.46	0.12	4.9	1.2
Mon R2 (1)	0.37	-0.18	3.0	2.0
(2)	0.23	-0.23	3.2	1.8
(3)	0.29	0.14	3.3	1.8
L1228	0.14	-0.17	6.0	1.0
L1551	0.18	0.60	4.6	1.2
HH83	0.12	-0.39	4.2	1.3

FIG. 8.— The tails of the centroid velocity pdfs in Figures 6 and 7 are here shown on log-log axes in order to emphasise any power law components ($\propto v^n$) which may be present. On such axes, power law forms would appear as straight lines. The abscissa in each panel runs from one to six times the pdf standard deviation, σ (see Table 4). Solid lines and asterisks denote the high velocity pdf tails and dashed lines and squares denote the low velocity pdf tails. All pdf tails are normalised to unity at a velocity of one σ . Plot symbols (asterisks and squares) represent the histogram pdf estimator, while lines (solid and dashed) represent the adaptive kernel pdf estimator. The dotted line in the lower left corner of each panel illustrates a power law with a similar slope for comparison.

and the fine structure in Orion B, region 1c, all of which appear in the histogram and adaptive kernel estimators, but are not well-represented by the Johnson system. In general, however, the Johnson system provides a reasonable analytic approximation to the pdfs presented here, and the parameters listed in Table 3, together with the analytic form given by equation (1) can therefore be used by observers and modellers alike to roughly reproduce and compare with our results.

The velocity pdfs for the Mon R2 subregions and the high-velocity portion of Orion B, region 1b appear nearly parabolic on the log-linear axes shown, implying Gaussian forms, although the far tails for these regions still may suggest exponentials or power laws. However, most of the other pdfs appear distinctly non-Gaussian. Several of the pdf tails, particularly for regions 2, 3, and 4 in Orion B, as well as L1228, L1551, and HH83, instead suggest nearly exponential (straight lines on the log-linear axes of 4 and 5), or possibly power-law (straight lines on the log-log axes of Figure 8) forms. These results are consistent with the derived stretching exponents, which lie near 2 for the three Mon R2 regions, and near 1.1 ± 0.3 for most of the others (see Table 4 and the associated discussion). In no case do we find pdf tails which decay more rapidly than Gaussian ($\beta \lesssim 2$ for all regions).

It should be noted that the more Gaussian nature of the

Mon R2 centroid velocity pdfs is not associated with particularly high or particularly low Mach numbers relative to the nine other regions (Table 1). However, relative to regions 2, 3, and 4 in Orion B, which have a comparable size and mass scale, the Mon R2 regions do possess somewhat lower Mach numbers (Table 1) and somewhat higher Reynolds numbers (Table 5). Also, differences in the distribution and dynamical influence of young, massive stars in Orion B and Mon R2 may be significant (see Appendix A).

The pdf shapes can be described further using the kurtosis values listed in Table 4. The three Mon R2 regions are all characterised by a kurtosis of about 3, again suggesting nearly Gaussian statistics. However, with the exception of Orion B, region 1a, the other pdfs have larger kurtosis values, ranging from 4 to 6 (As explained above, the large kurtosis of 11 found for Orion B, region 2, is probably anomalous, due to overlapping with region 3). These results are in strong contrast to those for incompressible turbulence, where the velocity pdf is nearly Gaussian (see references given in §2.1), and they imply a fundamental difference between the physics of interstellar supersonic, driven turbulence and incompressible turbulence. An interpretation of the difference based on the fact that kinetic energy is not a conserved quantity in supersonic turbulence is discussed in Paper III.

Our results for molecular gas powered by internal sources are basically in agreement with the velocity pdfs from optical line and HI emission and absorption components for cool atomic gas displayed in Fig. 1. Since these latter regions are generally not self-gravitating and contain no internal power sources, the similarity may imply that the non-Gaussian behaviour is either a result of nonlinear advection or that the latter regions possess turbulence that is powered by external sources.

However, we note that at least 3 of the 11 subregions studied here *do* appear Gaussian, so our results should not be taken as an indication that all molecular clouds have exponential or even non-Gaussian velocity distributions. However since the 3 regions in question are all in the same GMC (Mon R2), we can say that the Gaussian distributions are definitely in the minority. A similar statement can be made for the optical and HI line results discussed in sec.2.2. Perhaps the most interesting challenge, theoretically, is to understand why the pdf is approximately exponential in most of the regions but apparently Gaussian in Mon R2.

4.2. Centroid Velocity Differences

As mentioned in §2.1, probability density functions of velocity derivatives or differences have proven to be an important and well-studied flow diagnostic in the context of incompressible turbulence, where they exhibit significantly non-Gaussian shapes. It is therefore of great interest to construct maps of centroid velocity differences in molecular clouds and investigate their statistical properties. In this section we apply such an analysis to our data sets.

The velocity differences we consider are defined as follows;

$$\Delta v(\mathbf{x}, \boldsymbol{\tau}) = v(\mathbf{x}) - v(\mathbf{x} + \boldsymbol{\tau}), \quad (6)$$

where $\mathbf{x}$ and $\boldsymbol{\tau}$ are two-element vectors corresponding to position on the plane of the sky and v is the centroid velocity. The displacement $\boldsymbol{\tau}$ is referred to as the two-dimensional vector lag. The structure function is just the mean square of these velocity differences, averaged over the area of the cloud and normalised with respect to the centroid velocity variance (e.g. MB). In other words, the structure function at a particular lag is the normalised variance of the difference pdf corresponding to that lag. It therefore represents only part of the information available from the full pdf, which we consider below.

Grey-scale images of the centroid velocity differences for each region are presented in Figures 9 and 10, for a fixed vector lag, in pixels, of $\boldsymbol{\tau} = [1, 1]$. In other words, the images of Figures 9 and 10 were obtained by displacing the maps of Figures 4 and 5 horizontally and vertically 1 pixel and then computing the difference as expressed by equation (6). Of particular interest when considering the centroid velocity differences is the spatial distribution of large-amplitude velocity differences—those events that compose the pdf tails. Such events have been emphasised in Figures 9 and 10 by using a 3-level grey scale. As indicated by the intensity table in Figure 10, all pixels in which the magnitude of the velocity difference is more than twice the standard deviation of the map are shown as white. Magnitude values between one and two standard deviations are shown as light grey, while values less than

one standard deviation are shown as dark grey.

It should be noted that many of the extreme events found in Figures 9 and 10 are likely spurious, influenced by low signal-to-noise ratios near the cloud edges and by residual emission from overlapping cloud components (see Appendix B). This can be seen especially in Orion B, region 1b, L1551, and in the easternmost portions of Orion B, regions 2 and 4. However, the distribution of extreme events within the main volume of the clouds is more reliable and generally reveals a very intermittent structure. In most cases, high-amplitude velocity differences appear to be well distributed throughout the clouds, although there is also some indication that some such events are correlated with the dense, star-forming cores of Orion B, region 4 and Mon R2, region 1 (see also Figs. 2 and 3). At larger lags, the distribution of large-amplitude events becomes less intermittent, as demonstrated in Figure 11 for Orion B, region 4.

An important question is whether or not the largest velocity differences occur in correlated, filamentary structures, as in the simulations of mildly supersonic, decaying turbulence discussed by Lis et al. (1996). We see little evidence for such structures in the images of Figures 9 and 10, which appear to have a relatively more “spotty” spatial distribution. Although noise-induced fluctuations near low-intensity cloud edges probably do contribute, the relatively spotty appearance persists even in the cloud interiors and near those cloud edges (e.g. the intensity ridge in the western portion of Orion B, region 4) where the intensity is high and the influence of noise is small (see Appendix B).

The qualitatively different spatial distribution of these observed velocity differences is likely due to the fact that the regions discussed here are highly supersonic and are continually driven by stellar energy input. In this case the velocity field is expected to be dominated by the compressional (shock) modes, not the vortical modes that dominated the simulations discussed by Lis et al. It is important to recognize that, except perhaps for scales smaller than about 0.1 pc, the turbulent velocity field of the ISM should be highly supersonic, and driven by shock interactions (Kornreich & Scalo 1998). Our results for the velocity differences, along with the velocity pdfs themselves, as discussed in §4.1 above, therefore suggest a fundamental difference between interstellar turbulence and incompressible turbulence, except perhaps in regions that have avoided energy input for a time long enough that they approach incompressible conditions by means of turbulent decay.

However Miville-Deschenes, Joncas, & Falgarone (1998) report, based on their study of HI in the Ursa Major cirrus cloud, that the spatial positions corresponding to the non-Gaussian tails of the difference pdf are primarily located along three filaments in the cloud, although differences images like our Figures 9 and 10 are not shown. Such an image of velocity differences *is* presented in the CO (2–1) study of the ρ Oph cloud by Lis, Keene, et al. (1998). Since the ρ Oph core is a region with active internal star formation, it should be more directly comparable with the regions studied in the present paper. The image of large velocity differences (Fig. 8 in Lis, Keene, et al. 1998) does indicate that some of the highest velocity differences are filamentary, although others are not. Thus the situation

FIG. 9.— Shown are grey-scale images of the centroid velocity differences for each subregion in Orion B, corresponding to a lag in pixels of $\tau = [1, 1]$. As indicated by the legend in Figure 10, the grey scale has only three shades, chosen to highlight the spatial distribution of large-amplitude difference values (white).

FIG. 10.— Similar to Figure 9, but for the Mon R2, L1228, L1551, and HH83 data sets. Again, all maps correspond to a lag in pixels of $\tau = [1, 1]$. The grey-scale intensity levels are based on the standard deviation of each map, as indicated by the legend.

FIG. 11.— Shown are grey-scale images of the centroid velocity differences for Orion B, region 4 at the lags indicated (measured in pixels). The dashed boxes indicate the displacement as specified by the lag vector, τ . The sense of the grey-scale is the same as in Figures 9 and 10.

is not clear-cut, and the most we can claim is that, for the regions studied here, there is little evidence for filamentary clustering of regions with large velocity differences. Quantification of this statement is beyond the scope of this paper, since establishing the reality or absence of filamentary structure is a difficult and unsolved problem, as is well-known from studies of the large-scale distribution of galaxies.

As in §4.1.1, we can investigate the velocity differences further by considering their probability density functions. In order to facilitate visualisation and improve statistics, we combine data from all lag angles to obtain pdfs as a function of the lag magnitude, $\tau \equiv |\tau|$. For each scalar lag, τ , measured in pixels, there is an associated pdf which describes the distribution of velocity difference values. The pdf variation with τ is illustrated in Figures 12 and 13. For each region, solid, dashed, and dotted lines correspond to scalar lags (τ) of 1, 5, and 20 pixels. All pdfs have been computed using the adaptive kernel estimator described in §4.1.1.

Some of the difference pdfs in Figs. 12 and 13 appear to change from nearly-exponential at small lags to more nearly-Gaussian at large lags, a behaviour which is similar to incompressible turbulence, the mildly supersonic decay simulations of Lis et al. (1996), and the observational results of Lis, Keene, et al. (1998) and Miville-Deschenes, Joncas, & Falgarone (1998). However many of the regions show a persistence of exponential behaviour even at large lags ($|\tau| = 20$ pixels), although sometimes with a cutoff at very large velocity differences. This persistence of exponential behaviour at large lags is seen in most of the observed regions, and again points to a difference between interstellar turbulence and incompressible turbulence. Alternatively, the absence of a clear transition to Gaussian

behaviour may simply reflect the fact that, at the largest lags examined, the velocity field is still correlated while such larger-scale correlations do not occur in the incompressible regime or in the regions examined by Lis, Keene, et al. (1998) and Miville-Deschenes, Joncas, & Falgarone (1998).

The increase of the width of the velocity difference pdf with increasing lag is qualitatively similar to results found for incompressible turbulence. However the same behaviour may be expected for any stochastic field: at larger separations, the variables become less correlated, and hence a larger fraction of the differences have large values. Basically this result only shows that the structure function (variance of velocity difference) increases with increasing lag, a property shared by a large class of stochastic processes (e.g. Burgers turbulence, which is entirely compressible). In order to constrain models for interstellar turbulence, it is the functional form of this increase, as well as the behaviour of other moments, which is crucial.

We therefore consider the variation with lag of the central sample moments of the difference pdfs, defined as in equations (2)–(5). The second moment, or variance, for each region is presented in Figure 14, as a function of the scalar lag, τ , expressed in parsecs. As mentioned above, the variance of the difference pdfs is simply the structure function, without the customary normalisation factor. The six subregions of Orion B are shown in frame (a), and frame (b) shows the three subregions of Mon R2 together with the L1228, L1551, and HH83 results. The variation of the normalised fourth moment, or kurtosis [eqn. (5)], with scalar lag for each region is displayed in Figure 15.

It is clear from Figures 14 and 15 that the variation of the difference moments over some lag ranges can be approximately described by a power-law dependence, which

FIG. 12.— Probability density functions of centroid velocity differences for the six subregions in Orion B. In each plot, solid lines correspond to a scalar lag, τ , of 1 pixel, while dashed and dotted lines correspond to scalar lags of 5 and 20 pixels respectively. All pdfs shown are computed using the adaptive kernel estimator described in §4.1.1.

FIG. 13.— Similar to Figure 12, but for the Mon R2, L1228, L1551, and HH83 data sets. Solid, dashed, and dotted lines again correspond to the velocity difference pdfs at scalar lags of 1, 5, and 20 pixels.

appears linear on the log-log axes shown. A characteristic power-law index can be derived for each region by measuring the mean logarithmic slope of each curve over some chosen range in lag. However, the choice of the fitting range is not straightforward—most of the curves in Figures 14 and 15 exhibit steep slopes at small lags which flatten out as the lag increases. This “flattening out” is a common feature of all stochastic fields, which approach a constant variance and kurtosis at lags far exceeding the field’s characteristic correlation length, or as in the present context, the effective width of the applied spatial filter. The fitting range should therefore not include large lags, where the slope is systematically small. On the other hand, sampling effects can influence the slope at the smallest lags (MB). We therefore chose the intermediate range of lags between 6 and 12 pixels (note that this corresponds to a different physical scale in each region), and applied power-law fits of the form:

$$\text{variance} \propto \tau^{\alpha_2} \quad \text{and} \quad \text{kurtosis} \propto \tau^{\alpha_4} . \quad (7)$$

The results yielded α_2 values between 0.64 and 1.05 (except for Orion B, region 1b, in which $\alpha_2 = 0.33$) and α_4

values between -0.11 and -0.88, with typical values of 0.85 and -0.5 respectively. These typical values are shown on the plots of Figures 14 and 15 for comparison with the data. We emphasise that these best-fit power law exponents depend significantly on the chosen fit range and should therefore be viewed only as a rough estimate of the logarithmic slope of the moment curves at intermediate lags. With the exception of the L1228 variance ($\alpha_2 = 0.66$), the smaller-scale, low-mass star-forming regions L1228, L1551, and HH83 exhibit somewhat steeper logarithmic slopes ($1.01 \leq \alpha_2 \leq 1.05$, $-0.88 \leq \alpha_4 \leq -0.64$) than the GMC regions ($0.33 \leq \alpha_2 \leq 0.90$, $-0.52 \leq \alpha_4 \leq -0.11$), possibly reflecting the influence of molecular outflows. The smallest slopes occur in the “cometary clouds”, regions 1a and 1b of Orion B, although it should be noted that these regions have the poorest statistics (in terms of both signal-to-noise and number of spectra).

The variance results reported here are consistent with the more detailed structure function analysis of MB. The reader is referred to that paper for further work and for a comparison with structure function indices predicted by various analytic and numerical turbulence models and

FIG. 14.— The variance of the difference pdfs is shown as a function of scalar lag, τ , measured in parsecs. Frame (a) is comprised of the six subregions of Orion B and frame (b) includes the three subregions of Mon R2 and the remaining L1228, L1551, and HH83 data sets. The assumed distances to each region are as listed in Table 1. A power law of index 0.85 is shown in each plot for comparison.

FIG. 15.— Similar to Figure 14, but here the kurtosis of the difference pdfs is plotted as a function of scalar lag, τ . As indicated by the legend in Figure 14, frame (a) corresponds to Orion B, and frame (b) corresponds to Mon R2, L1228, L1551, and HH83. A power law of index -0.5 is shown in each plot for comparison.

with empirical scaling relations from interstellar observations (e.g. “Larson” scaling) and laboratory turbulence experiments (e.g. “Kolmogorov” scaling).

The variation of kurtosis (flatness) with lag can be compared with the incompressible turbulence experiments of VanAtta & Antonia (1980) and the multifractal models of Eggers & Wang (1998), which exhibit a basically constant kurtosis out to some critical lag (which depends on the Reynolds number), beyond which the kurtosis decreases with increasing lag. However, except for a narrow transition region near the critical lag, this decrease is much slower than that found in the present work.

We believe that the ability of hydrodynamical simulations to account for the observed variations of second (Fig. 14) and fourth (Fig. 15) moments of the velocity difference pdfs with lag provides a decisive test of turbulence models, although it must be remembered that the observations refer to regions that are supersonic and contain internal turbulent power sources.

4.3. Reynolds Numbers

The dynamical information available in the centroid velocity and velocity difference maps considered above can be used to assess the “degree” of turbulence in each region, as quantified by the Reynolds number. The Reynolds number $Re = v\ell/\nu$ is defined only with respect to the scale ℓ , which might be the size of the region, the correlation length (integral scale), or any other length that measures a useful characteristic spatial scale of the turbulence. One such scale which has proven especially useful in characterising incompressible turbulence experiments and simulations is the Taylor microscale, which measures the average spatial extent of velocity gradients. For incompressible turbulence, these gradients are entirely due to the vorticity, so the Taylor microscale is defined as (e.g. Lesieur 1990, p. 144)

$$\ell_T = \frac{u_{rms}}{\langle (\nabla \times \mathbf{u})^2 \rangle^{1/2}} \quad (8)$$

where u_{rms} is the root-mean-square fluid velocity. For compressible turbulence the velocity gradients are due to a combination of vortical (rotational, solenoidal) and compressible (irrotational, dilatational) modes. For highly supersonic turbulence the vorticity in eqn. (8) might be replaced by the dilatation $\nabla \cdot \mathbf{u}$. (In this case the simple scaling between Taylor scale Reynolds number and integral scale Reynolds number [eq. VI-6-7 in Lesieur 1990, p. 144] is lost because there are no relations between enstrophy and viscous dissipation rate and the dissipation rate cannot be taken as u^3/ℓ as for the dissipationless Kolmogorov energy cascade.)

The Taylor scale Reynolds number is generally smaller than the integral scale Reynolds number. For reference, simulations of incompressible turbulence are only capable of generating turbulence with Taylor scale Reynolds numbers of at most $10^2 - 10^3$, while experiments reach Taylor scale Reynolds number of at most $\sim 10^3 - 10^4$ (see, for example, Kailasnath, Sreenivasan, & Stolovitzky 1992, She et al. 1993, and references therein).

For empirical studies such as the present one, it is appropriate to use the rms value of the measured velocity gradient in the denominator of eqn. (8). This is clearly an

important quantity, since it gives a measure of the characteristic scale over which turbulent interactions of any sort (e.g. shocks, vorticity stretching) occur. A problem is that estimation of centroid velocity gradients across the line of sight will be strongly amplified by uncertainties in the centroid velocities, while if we try to avoid this problem by smearing the centroid velocity field with a filter, the resulting rms velocity gradient will decrease with increasing filter size. A convenient way to circumvent these problems is to take the appropriate mean velocity gradient as the square root of the variance of the velocity difference pdf taken at the smallest separation (one pixel), which we will call σ_s , divided by the physical length corresponding to this separation, which we will call s . The reasoning is that the average taken when computing the variance of the difference pdf will greatly reduce the noise due to uncertainties in individual centroid velocities. We therefore define the Taylor scale as

$$\ell_T = \frac{\sigma_c s}{\sigma_s} \quad (9)$$

where σ_c is the rms centroid velocity fluctuation amplitude. Taking the velocity occurring in the Reynolds number as σ_c , this gives a Taylor scale Reynolds number

$$Re_T = \frac{\sigma_c^2 s}{\nu \sigma_s} \quad (10)$$

We take σ_c to be the dispersion of the mean spectral line profile for each region, given by the “parent dispersion” σ_p in Table 2 of MB. Alternatively, the “turbulent dispersion”, σ_t in Table 2 of MB, could be used, but it includes uncertainties due to instrumental noise and it removes the influence of large-scale velocity variations under the assumption that they are uncorrelated with small-scale fluctuations. Since large scales in turbulent flows typically contain much of the kinetic energy and play an important role in the nonlinear energy transfer among modes, σ_p is the most appropriate dispersion measure to use in computing an effective Reynolds number. However, the difference is relatively minor.

Note that our definition of the Taylor scale (eqn. 9) and the associated Reynolds number (eqn. 10) depend on the scale at which they are computed—i.e. the spatial resolution of the map. However, this is reasonable because it is the scale of the fluid motions we are sampling - we have no information about the dynamics on scales smaller than the resolution or larger than the emission region. The values may change if higher resolution maps become available, but only if there is a large amount of power at high spatial frequencies. The range of values found for the Taylor scale, 3 to 10 times larger than the grid spacing, suggests that this may not be the case. Projection effects and line-of-sight averaging also influence the computation of the Taylor microscale, but these effects are very difficult to quantify without some assumption about the three-dimensional nature of the velocity field. They could lead to an overestimate of the Taylor microscale and the corresponding Reynolds number, although to our knowledge this has never been demonstrated.

The molecular viscosity is estimated using a mean free path approximation, $\nu = v_{th}/\sigma n$, where v_{th} is the rms thermal velocity $(kT/\mu m_H)^{1/2}$, n is the particle number

TABLE 5
REYNOLDS NUMBERS AND TAYLOR SCALES

Region	L (pc)	n (10^2 cm^{-3})	σ_c (km s^{-1})	s (pc)	σ_s (km s^{-1})	$Re_L/10^7$	$Re_T/10^6$	ℓ_T (pc)
Orion B (1a)	6.0	2	0.55	0.12	0.10	1.2	1.3	0.66
(1b)	8.6	2	0.55	0.12	0.19	1.7	0.69	0.35
(1c)	17	2	0.83	0.12	0.21	5.1	1.4	0.47
(2)	30	2	1.15	0.12	0.21	12	2.7	0.66
(3)	33	2	2.05	0.12	0.41	24	4.4	0.60
(4)	40	2	1.31	0.12	0.25	19	3.0	0.63
Mon R2 (1)	41	6	1.12	0.24	0.19	50	17	1.4
(2)	32	6	0.74	0.24	0.12	26	12	1.5
(3)	50	6	0.83	0.24	0.16	45	11	1.2
L1228	2.6	10	0.74	0.087	0.064	3.5	13	0.59
L1551	1.8	10	0.50	0.041	0.085	1.6	2.2	0.24
HH83	0.67	25	0.42	0.014	0.042	1.2	2.6	0.14

density, and σ is the collisional cross section. We adopt the cross section for $H_2 - H_2$ elastic collisions as about 10^{-15} cm^2 . It is unclear to us whether the absence of a dipole moment for the H_2 molecular should alter the elastic cross section compared to typical values for other atoms and molecules reported in the literature; we assume it does not. The particular densities are taken as the rough estimates given by MB: $n(\text{cm}^{-3}) = 2500$ for HH83, 1000 for L1228 and L1551, 600 for Mon R2, and 200 for the Orion B subregions. We adopted a characteristic temperature of 20 K for all the regions.

Then the Taylor scale Reynolds number can be expressed as

$$Re_T = 1.8 \times 10^6 n_2 \sigma_{c,5}^2 s_{pc} / \sigma_{s,5} \quad (11)$$

where $n_2 = n/100 \text{ cm}^{-3}$, $\sigma_{c,5}$ is the turbulent velocity dispersion in km s^{-1} , s_{pc} is the linear scale corresponding to a one pixel lag at the adopted distance of the region, and $\sigma_{s,5}$ is the standard deviation of the velocity difference pdf at this lag, in km s^{-1} . For comparison, the Reynolds number corresponding to the scale L of the observation, taken as the geometrical mean of the semiminor and semimajor axis of the emitting region on the plane of the sky, is

$$Re_L = 1.8 \times 10^6 n_2 \sigma_{c,5} L_{pc} . \quad (12)$$

We computed these dimensionless numbers for each region and the results are given in Table 5. The region scale Reynolds number Re_L is, as expected, very large, varying between about 1×10^7 to 5×10^8 . This range is very nearly the same as the estimates of Re_L given by Myers & Khersonsky (1995) for a sample of “diffuse,” “dark,” and “giant” clouds. Surprisingly, the Taylor scale Reynolds numbers, while generally smaller than Re_L , are still very large, between 7×10^5 and 2×10^7 . As mentioned above, the Taylor microscales, which measure the rms scale of velocity gradients, are, in units of the resolution s , $\sigma_c/\sigma_s = 3$ to 10, suggesting that we are resolving the rms gradient scale. However, the very large values of the Taylor scale Reynolds

number mean that, at the smallest resolvable scales, the advection term in the momentum equation, as measured by the velocity gradients, is still huge compared to viscous dissipation, implying that we have not yet resolved the dissipation scale, so velocity gradients must still be present on spatial scales smaller than the sampling grid of the observations. Considering the numerical diffusion/artificial viscosity in existing numerical codes, simulations are far from approaching the resolution necessary to realistically represent interstellar turbulence. However, in the presence of a magnetic field, the dissipation might be due to ion-neutral friction or magnetic reconnection rather than ordinary viscosity (Myers & Khersonsky 1995), making the situation less severe.

A surprising result apparent in Table 5 is that the Taylor scale ℓ_T is approximately constant among most of the subregions of Orion B ($\ell_T \approx 0.6 \text{ pc}$) and of Mon R2 ($\ell_T \approx 1.4 \text{ pc}$). The difference between Orion B and Mon R2 might be attributed to the differing spatial resolutions, since ℓ_T (eqn. 9) is proportional to the size of a resolution element, and Mon R2 is roughly twice as distant as Orion B. The same distance scaling is apparent for L1551, and qualitatively for L1228 and HH83, where ℓ_T is only 0.14 pc: HH83 is at about the same distance as Orion B, but was observed at about eight times better resolution. These results may suggest that the Taylor microscale is approximately constant in all the regions observed when the reference spatial scale s (eqn. 9) is normalised to the same value. This is a particularly intriguing result, since it would imply that the characteristic rms scale of velocity gradients, measured relative to the rms global velocity field, is a characteristic length scale independent of differing physical conditions in the regions studied. However, we are unable to assign a value to this scale because it depends on the adopted scale over which the gradients are measured, taken here as the size of a resolution element.

Experiments (VanAtta & Antonia 1980; Tabeling et al. 1996) and multifractal models (Biferale 1993; Eggers & Wang 1998) of incompressible turbulence indicate that the kurtosis (flatness) of the velocity difference pdf should

increase with Re_L as about $R_L^{0.15}$, but for Taylor scale Reynolds numbers greater than about 700 the kurtosis decreases with Re_L . We find no evidence for such behaviour in the observations presented here, again suggesting that highly compressible turbulence differs significantly for incompressible turbulence. However it should be noted that both Re_L and Re_T are proportional to adopted average densities, which are very uncertain, and so errors in the density estimates may mask any correlations.

We have not attempted to estimate magnetic Reynolds number (see Myers & Khersonsky 1995) for each region, because the magnetic viscosity depends on the magnetic field strength and the ionized fraction, quantities which are unknown for the present regions. The significance of these numbers for the propagation of MHD waves is discussed in Myers & Khersonsky (1995).

5. SUMMARY

The probability distribution function (pdf) for fluctuations of molecular line centroid velocities, and line centroid velocity differences at different separations, have been estimated for a number of local regions with active internal star formation. The data consist of a total of over 75,000 ^{13}CO line profiles covering five different molecular clouds. The internal stellar power sources include only low-mass protostellar winds in three regions (L1228, L1551, HH83) but extend to massive stars in the remaining two regions (Orion B and Mon R2). The GMCs Orion B and Mon R2 were subdivided into six and three kinematically distinct subregions respectively in order to isolate particular cloud components. The total sample therefore numbers twelve distinct regions, each composed of about 1000 to 25000 independent spectra. Centroid velocity fluctuation maps were constructed by interpolating these spectra onto regular spatial grids and then applying spatial filters in order to remove large-scale gradients. These maps are displayed as images in Figs. 4 and 5.

The pdf of centroid velocities for each region was estimated using the classical histogram, a non-parametric adaptive kernel estimator, and the parametric Johnson estimator. The methods generally agree well except in the far tails of the pdfs. Although the Mon R2 regions exhibit nearly-Gaussian pdfs, except possibly in the far tails, all the other regions show strong excesses relative to a Gaussian, often suggesting nearly-exponential or power-law forms. Sample moments and stretched exponential fits to the pdfs were presented as quantitative measures of the departure from Gaussian statistics. These results confirm and extend the general conclusions reached by MS.

Centroid velocity pdfs for diffuse interstellar HI regions, constructed from older published optical line and HI emission and absorption line data, are also presented, and also show strong evidence for non-Gaussian, nearly exponential, pdfs. The similarity with most of the molecular regions studied here suggests that either the exponential pdfs are not a product of stellar activity or that the internal velocity fields of the diffuse regions have been strongly affected by disturbances from external sources. These exponential centroid pdfs do not agree with the nearly-Gaussian pdfs found in numerical simulations of freely-decaying marginally supersonic turbulence by Lis et

al. (1996). The pdfs are also markedly different from the nearly Gaussian velocity pdfs found in studies of incompressible turbulence. A theoretical interpretation of the centroid pdfs is presented in a separate paper (Paper III).

We also constructed the pdfs of centroid velocity differences for lines of sight separated by different scalar spatial lags. In agreement with other recent observational work on this function, and with studies of incompressible turbulence, we find nearly exponential difference pdfs at small lags which broaden with increasing lag. However, spatial images of the centroid differences show a “spotty” distribution for the largest velocity differences, with little evidence for the filamentary structures predicted by simulations of decaying marginally supersonic turbulence—structures which should be due to vorticity.

We used the velocity difference pdfs to estimate the Taylor microscale, which is the rms scale of velocity gradients. The Taylor microscale Reynolds number for all the regions is very large, $\sim 10^5$ to 10^6 , indicating that even at the smallest resolvable scales, the advection term in the momentum equation is still huge compared to viscous dissipation.

Our observational results suggest fundamental differences between turbulence in both star-forming regions and diffuse atomic interstellar clouds as compared with incompressible or mildly-supersonic decaying turbulence. These interstellar gas motions are characterised by very large Reynolds numbers, a high degree of compressibility, continuous energy and momentum injection by internal and external power sources, and various sources of anisotropy, including rotation, shear, magnetic fields and self-gravity. It might be necessary to incorporate some or all of these characteristics into numerical and theoretical models before full agreement with observations can be achieved. The complexity of interstellar turbulence suggests that it may exhibit some degree of nonlinear self-organisation, producing coherent structures which could contribute non-Gaussian components to the velocity and velocity difference pdfs. Such behaviour has recently been demonstrated in three-dimensional numerical simulations of turbulent incompressible shear flow (Pumir 1996; Lamballaise, Lesieur, & Matais 1997) and turbulent compressible convection (Brandenburg et al. 1996). On the other hand, it may be that some rather simple and fundamental property of the ISM is behind its statistical behaviour. For example, stellar winds, outflows, and superbubbles can all produce non-Gaussian velocity distributions with power-law pdf tails (e.g. Silk 1995; Oey & Clarke 1998). Alternatively, as another example, the ISM is a system in which kinetic energy is not a global invariant conserved by the advection operator as it is for incompressible turbulence; in such a situation it is possible to obtain exponential velocity distributions, as arise in simulations of systems of interacting wind-driven shells (Chappell & Scalo 1999). In Paper III, we will further address these and related issues regarding the theoretical implications and interpretation of the results reported in the present paper.

This work was supported by NASA grant 5-3107 to JS, and by the National Research Council, through a postdoctoral fellowship to MM.

APPENDIX

APPENDIX A: OVERVIEW OF REGIONS STUDIED

All of the regions considered here are actively forming stars and there is evidence to suggest that the stars which have been produced have played a significant role on the dynamics of their parent clouds. In this appendix we provide a brief description of the environments and physical conditions in each of the regions studied in order to aid in the physical interpretation of the statistical analysis of §4

Orion B

The Orion giant molecular cloud (GMC) complex is one of the nearest and most popular sites for studying massive star formation and its interaction with surrounding interstellar material. The GMC is comprised primarily of the A cloud (L1641), which lies behind the Trapezium stellar cluster and the well-known Orion nebula, and the B cloud (L1630), which is of a comparable size and mass and extends out to the northeast. The molecular gas in the region has been mapped extensively in ^{12}CO by Maddalena et al. (1986), in the associated dust emission using IRAS (Beichman 1988; Robinson 1984; Wood, Myers, & Daugherty 1994), and in other isotopes of CO and CS (e.g. Bally et al. 1991; Lada, Bally, & Stark 1991; Kramer, Stutzki, & Winnewisser 1996). The star formation activity and associated cloud dynamics have been reviewed by, e.g., Genzel & Stutzki (1989).

The Orion GMC complex may have originated through the fragmentation and local gravitational collapse of the expanding Gould's Belt supershell about 10 to 20 Myr ago Bally (1995) or from the collision of an infalling high-latitude cloud with the galactic plane (Franco et al. 1988). Massive star formation began by 12 Myr ago and produced the I Orion OB association. There is good evidence that these young, hot stars have had a profound influence on their parent clouds by accelerating, ionizing, ablating, compressing, and disrupting them through supernovae, radiation, and stellar winds (Bally, Langer, Stark, & Wilson 1987; Bally, Stark, Wilson, & Langer 1989; Bally et al. 1991; Genzel & Stutzki 1989). The large-scale velocity gradients and morphologies of the main Orion A and B clouds as well as the “wind-swept” or “cometary” appearance of a number of nearby smaller clouds, with “tails” directed away from the OB association (as in regions 1a and 1b in the present study), provide a strong indication that the association has played an important role in their dynamics.

The Orion B mapping considered here (Fig. 2) covers a number of regions which are still actively forming stars. The reflection nebulae NGC 2068 and 2071 are located near emission peaks in what we have defined as region 2, and the main peaks along the emission ridge in region 4 are near the well-known HII regions NGC 2023 and 2024 and the Horsehead Nebula B33. Other notable objects within the mapped region include a large number of reflection nebulae, embedded infrared sources, HII regions, Herbig-Haro objects, H_2O masers, and molecular outflows (see Maddalena et al. 1986, Reipurth 1985, Lada, Bally, & Stark 1992, Chandler & Carlstrom 1996 and references therein). In an infrared survey of the area, Lada, DePoy, et al. (1991) identified approximately 1000 sources, about half of which are probably embedded in the Orion B cloud. Most of these sources are clustered, with ~ 330 near NGC 2023 and 2024 in our region 4, and another ~ 300 near NGC 2068 and 2071 in our region 2. Active star formation is even occurring in at least some of the smaller clouds in region 1, as demonstrated by Reipurth & Olberg (1991), who report several Herbig-Haro jets and molecular outflows in L1617 (which we have labelled region 1b).

In summary, the Orion B observations studied here represent a very dynamic region where the molecular gas is substantially influenced by supernovae, radiation, and winds from the massive stars in the nearby OB association and from embedded young stars which are continually forming within.

Mon R2

The Monoceros R2 molecular cloud is comparable in size and CO luminosity to Orion B. The large-scale distribution of ^{12}CO follows well the area spanned by the Lynds dark clouds L1643, L1644, L1645, and L1646 (Maddalena et al. 1986).

The core of the Mon R2 cloud lies at the origin of the coordinate system used in Figure 3a, near the predominant CO emission peak, and is a well-studied region of ongoing high-mass star formation. It has been extensively mapped in CO isotopes, CS, HCN, H_2CO , NH_3 , and HCO^+ , and is associated with a number of B stars and reflection nebulae, a large cluster of embedded infrared sources, a compact HII region, enhanced X-ray emission, and both H_2O and OH masers (see Beckwith et al. 1976, Thronson et al. 1980, Montalbán et al. 1990, Torrelles et al. 1990, Gonatas, Palmer, & Novak 1992, Giannakopoulou et al. 1997, Tafalla et al. 1997, Gregorio et al. 1998, and references therein). The embedded cluster has recently been studied in detail by Carpenter et al. (1997), who estimate that its population numbers at least 475 stars. Also associated with this core region is one of the largest, most massive molecular outflows known, extending for at least 4 parsecs along its axis and involving almost $200 M_\odot$ of interstellar material (Wolf, Lada, & Bally 1990; Meyers-Rice & Lada 1991; Xie, Goldsmith, & Patel 1993). The outflow is bipolar, centered near the middle of the embedded cluster, and dominates the dynamics of the molecular cloud within at least several arcminutes of the core (corresponding to the origin in Figure 3a). A separate outflow, located about $75''$ from the main embedded cluster, was recently identified by Tafalla et al. (1997).

A second prominent CO emission peak lies about $45'$ to the east (in region 2), and corresponds to the molecular core known as GGD12-15 (Little, Heaton, & Dent 1990). Evidence indicates that this region too is actively forming stars and is also associated with a large bipolar molecular outflow, a compact HII region, several infrared sources, and an H_2O maser (Rodríguez et al. 1980; Olofsson & Koornneef 1985; Little, Heaton, & Dent 1990). Furthermore, there are a

number of IRAS point sources tracing the sharp emission ridges in regions 2 and 3 as well as additional infrared sources and reflection nebulae scattered primarily throughout regions 1 and 2 and another HII region near the northeastern edge of region 1 (Xie & Goldsmith 1994). At least 30 to 40 separate infrared and X-ray sources have been identified outside the main Mon R2 core (Xie & Goldsmith 1994; Gregorio et al. 1998).

Loren (1977) studied the large-scale kinematics of the region using CO, H₂CO, and near-infrared observations and he interpreted velocity gradients across the cloud as a combination of rotation and collapse. However, the more recent, higher-resolution mappings by Xie & Goldsmith (1994) have revealed a more complicated intensity and velocity structure, characterised by sharp emission ridges in the western and northeastern portions of the cloud and a southeastern portion with a lower LSR velocity and a more “wispy” appearance. This overall structure is confirmed by the observations presented here and is the basis for the region decomposition described in §3. Xie & Goldsmith (1994) interpreted the relative blueshift of region 1 as evidence for a large expanding shell, involving $\sim 4 \times 10^4 M_\odot$ of material and moving toward us with a velocity of $\sim 3\text{--}4 \text{ km s}^{-1}$. Fairly recent, large-scale dynamical events are also suggested by the sharpness of the emission ridges in regions 2 and 3 and the evidence for associated star formation (see the preceding paragraph), which may indicate shock compression of the molecular gas.

In summary, Mon R2 is a giant molecular cloud comparable in scale to Orion B. Although there is no analogous OB association disrupting the molecular gas to the same degree as in Orion, active star formation is indeed occurring within Mon R2 and there is also some evidence suggesting violent dynamical events on the scale of the cloud.

L1228 and L1551

The Lynds dark clouds L1228 and L1551 are best known for the molecular outflows found within them. Both regions are nearby, well-studied sites of ongoing low-mass star formation.

The L1228 cloud is located in the Cepheus flare, in a ring of molecular gas which seems to be part of a 4×10^4 year old supernova remnant (Grenier et al. 1989; Yonekura et al. 1997). At the center of the coordinate system in Figure 3b lies the young stellar object IRAS 20582+7724 and associated with it, extending northeast and southwest, is a large ($18' \times 9'$), well-collimated, bipolar molecular outflow (Haikala & Laureijs 1989; Bally et al. 1995; Anglada, Sepulveda, & Gomez 1997; Tafalla & Myers 1997). The orientation and sense of the outflow axis is roughly the same as the large-scale velocity gradient apparent in Figure 3b. Also, Bally et al. (1995) have argued that there is at least one other YSO producing an outflow in the L1228 cloud core and have reported a number of HH objects in the vicinity. Further evidence for active low-mass star formation has been provided by Ogura & Sato (1990), who identified 69 H α emission stars in the area, nine of which are concentrated near the L1228 cloud core.

L1551 occupies a small area along the edge of the Taurus-Auriga molecular cloud complex (e.g. Ungerechts & Thaddeus 1987) and is one of the most fertile centers of low-mass star formation known. The large ($10' \times 35'$), collimated, bipolar molecular outflow in the southeastern portion of the cloud associated with the infrared source IRS 5 (which lies at the origin of the coordinate system in Fig. 3c) is one of the archetypical examples of the outflow phenomenon in young stellar objects (YSOs) and has been the subject of much observational and theoretical research (see Snell, Loren, & Plambeck 1980, Cabrit & Bertout 1986, Uchida et al. 1987, Moriarty-Schieven & Snell 1988, Stocke et al. 1988, Bachiller, Tafalla, & Cernicharo 1994, Davis et al. 1995, and references therein). In the coordinate system chosen in Figure 3c, the outflow axis is oriented approximately horizontally, with blue-shifted gas toward the right of the origin and red-shifted gas toward the left, and it occupies much of the lower portion of the map. Also associated with IRS 5 and its accompanying outflow are a large reflection nebula known as HH102 and several other Herbig-Haro objects, including HH28 and HH29 (e.g. Graham & Rubin 1992).

In addition to the IRS 5 vicinity, there are several other regions throughout the L1551 cloud which exhibit strong evidence for ongoing low-mass star formation. Especially several arcminutes to the north, near the very young T Tauri stars HL and XZ Tau and the YSO HH30, a large number of Herbig-Haro objects, molecular outflows, optical jets, infrared sources, and excess H α emission regions have been identified (Mundt, Ray, & Buhrke 1988; Mundt et al. 1990; Graham & Heyer 1990; Pound & Bally 1991 1991). Recent X-ray observations by Carkner et al. (1996) and CCD and spectroscopic observations by Briceno et al. (1998) have increased the total number of known T Tauri stars in the L1551 cloud to at least 26, along with one young B9 star.

HH83

The structure, dynamics, and nature of the HH83 molecular cloud have been described in detail by Bally, Castets, & Duvert (1994). The innermost core has also been mapped in CS emission by Nakano et al. (1994). This cloud is smaller than the others in our study (see Table 1), and is located at the extreme western edge of the Orion A cloud (L1641), within an area to the northwest of NGC1999 which has a high concentration of Herbig-Haro objects and several CO outflows, signifying active star formation in the region.

An embedded star—the infrared source HH83 IR—lies in the southwestern part of the cloud, at the origin of the coordinate system shown in Figure 3d, and an associated optical jet extends out toward the northwest for at least $32''$, terminating in a conical bow-shock about $150''$ from the central IR source (e.g. Reipurth 1989). Also associated with the embedded star and the jet is a low-velocity ($\approx 5 \text{ km s}^{-1}$), poorly collimated, bipolar outflow (Bally, Castets, & Duvert 1994). The outflow does appear in molecular emission lines, but there is good evidence that it is in a late stage of its evolution, having “blown out” of its parent cloud, and is currently depositing most of its energy and momentum into the atomic inter-cloud medium. Although the ¹³CO-emitting gas does not trace the full extent of the outflow, small-scale velocity gradients near

the central IR source and evidence for evacuated cavities along the jet axis suggest that the outflow has had a substantial dynamical influence on the molecular gas in the vicinity.

The axis of the outflow and jet, like several others in the region, is approximately aligned with the mean, large-scale magnetic field (directed northwest/southeast). The cloud is elongated perpendicular to this direction and, at least in the northern portion, exhibits a velocity gradient which may indicate gravitational collapse along the mean magnetic field lines on a timescale of $\sim 3 \times 10^5$ years. Also, the large-scale gradient across the cloud may indicate rotation about an axis aligned with the magnetic field and outflow. Furthermore, the cloud's proximity to the Orion OB association to the north suggests that it may have been accelerated southward relative to the surrounding gas by ablation, supernovae, and stellar winds. The associated compression may have triggered gravitational collapse in the southern end of the cloud, and may account for why star formation is currently occurring in the southern portion, but seems to be absent in the northern portion.

APPENDIX B: SOURCES OF ERROR

The influence of instrumental noise on the centroid velocity fluctuations was studied by Miesch & Bally (1994), and we only summarise their results here. For each region studied in that (and in this) paper, they approximated the mean spectrum as a Gaussian with a peak, dispersion, and center determined respectively by the observed mean brightness temperature, linewidth, and LSR velocity. The rms brightness temperature fluctuation due to instrumental noise was then estimated for each spectrum based on the effective integration times and detector temperatures, and a mean rms noise fluctuation was computed for each data set. A series of simulations was then performed for each region, based on the integration window used for the observations, the model Gaussian mean line profiles, and the mean noise fluctuations. In the simulations, white noise of the appropriate rms amplitude was added to the model profiles and the centroid velocity was computed. This operation was repeated (typically about 10000 times) with different random number seeds for the white noise, and the variance of the centroid velocity in these simulations was computed. The results indicate that the noise-induced fluctuations are very small, typically increasing the standard deviation of the centroid velocity by only a few percent (compare σ_c^* to σ_c in Table 2 of Miesch & Bally 1994).

Although these simulations do not directly address the influence of noise on the far tails of the centroid velocity and centroid velocity difference pdfs, they do suggest that this influence is minor, except for those spectra near cloud edges where the brightness temperature drops substantially below the mean brightness temperature. As discussed in §3, threshold integrated intensity levels were used to eliminate spectra with very low signal-to-noise, but some questionable spectra near cloud edges still remain. For example, the extreme velocity and velocity difference values near the edges of Orion B region 1b (Figs. 4 and 9) and L1551 (Figs. 5 and 10) may be influenced by noise.

In some areas, the influence of kinematically distinct, spatially overlapping emission may also bias the centroid pdfs, although the velocity ranges were carefully chosen to minimise this effect (see §3). This is particularly the case for the eastern portion of Orion B, region 4 (Figs. 4 and 9).

REFERENCES

- Adams, W. S. 1949 ApJ, 109, 354
 Adams, F. C. and Wiseman, J. J. 1994, ApJ, 435, 693
 Alves, J., Lada, C. J., and Lada, E. A. 1998, ApJ, in press
 Anglada, G., Sepulveda, I., & Gomez, J. F. 1997, A&A Suppl. Ser., 121, 255
 Anselmet, F., Gagne, Y., Hopfinger, E. J., & Antonia, R. A. 1984, J. Fluid Mech., 140, 63
 Anthony-Twarog, B. J. 1982, AJ, 87, 1213
 Bachiller, R., Tafalla, M., & Cernicharo, J. 1994, ApJ, 425, L93
 Bally, J., Langer, W. D., Stark, A. A., & Wilson, R. W. 1987, ApJ, 312, L45
 Bally, J., Stark, A. A., Wilson, R. W., & Henkel, C. 1987b, ApJS, 65, 13
 Bally, J., Stark, A. A., Wilson, R. W., & Langer, W. D. 1989, in *The Physics and Chemistry of Interstellar Molecular Clouds*, ed. G. Winnewisser & J. T. Armstrong (Berlin: Springer-Verlag), 81
 Bally, J., Langer, W. D., Wilson, R. W., Stark, A. A., & Pound, M. W. 1991, in *Fragmentation of Molecular Clouds and Star Formation*, IAU Symp. no. 147, ed. E. Falgarone, F. Boulanger & G. Duvert (Dordrecht: Kluwer Academic Publishers), 11
 Bally, J., Castets, A., & Duvert, G. 1994, ApJ, 423, 310
 Bally, J., Devine, D., Fesen, R. A., & Lane A. P. 1995, ApJ, 454, 345
 Bally, J. 1995, in *The Physics and Chemistry of Interstellar Molecular Clouds*, ed. G. Winnewisser and G. C. Pelz, (Berlin: Springer-Verlag), 118
 Barranco, J. A. & Goodman, A. A. 1998, ApJ, 504, 207
 Batchelor, G. K. 1953, *The Theory of Homogeneous Turbulence*, (Cambridge: Cambridge University Press)
 Beichman, C. A. 1988, *Astrophys. Lett. & Commun.*, 27, 67
 Beckwith, S., Evans, N. J. II, Becklin, E. E., & Neugebauer, G. 1976, ApJ, 208, 390
 Bernardeau, F. 1994, ApJ, 433, 1
 Biferale, L. 1993, *Phys. Fluids A*, 5, 428
 Blaauw, A. 1952, *Bull. Astron. Inst. Netherlands*, 11, 459 (B52)
 Blitz, L., Magnani, L., & Wandel, A. 1988, ApJ, 331, L127
 Brandenburg, A., Jennings, R. L., Nordlund, Å, Rieutord, M., Stein, R. F., & Tuominen, I. 1996, *J. Fluid Mech.*, 306, 325
 Briceno, C., Hartmann, L., Stauffer, J., & Martin, E. 1998, AJ, 115, 2074
 Cabrit, S. & Bertout, C. 1986, ApJ, 307, 313
 Carkner, L., Feigelson, E. D., Koyama, K., Montmerle, T., & Reid, I. N. 1996, ApJ, 464, 286
 Carpenter, J. M., Meyer, M. R., Dougados, C., Strom, S. E., & Hillenbrand, L. A. 1997, AJ, 114, 198
 Castaing, B., Gagne, Y., & Hopfinger, E. J. 1990, *Physica D*, 46, 177
 Catelan, P., & Moscardini, L. 1994a, ApJ, 426, 14
 ———. 1994b, ApJ, 436, 5
 Chandler, C. J. & Carlstrom, J. E. 1996, AJ, 114, 198
 Chappell, D. & Scalo, J. M. 1998, ApJ, in press
 ———. 1999, in preparation
 Chen, H., Herring, J. R., Kerr, R. M., & Kraichnan, R. H. 1989, *Phys. Fluids A*, 1, 1844
 Chen, S., Doolen, G., Herring, J. R., & Kraichnan, R. H. 1993, *Phys. Rev. Lett.*, 70, 3051
 Crovisier, J. 1978, A&A, 70, 43 (C78)
 Davis, C. J., Mundt, R., Eisloffel, J., & Ray, T. P. 1995, AJ, 110, 766
 Dickey, J. M., Salpeter, E. E., & Terzian, Y. 1978, ApJS, 36, 77
 Dubinski, J., Narayan, R., & Phillips, T. G. 1995, ApJ, 448, 226
 Eggers, J. & Wang, Z. J. 1998, *Phys. Rev. E*, 57, 4281
 Elias, J. H. 1978, ApJ, 224, 857
 Falgarone, E. 1989, in *IAU Colloq. 120, Structure and Dynamics of the Interstellar Medium*, ed. G. Tenorio-Tagle, M. Moles, and J. Melnick (Berlin: Springer-Verlag), p.68
 Falgarone, E. & Lequeux, J. 1973, A&A, 25, 253
 Falgarone, E. & Phillips, T. G. 1990, ApJ, 359, 344
 ———. 1991, in *Fragmentation of Molecular Clouds and Star Formation*, IAU Symp. no. 147, ed. E. Falgarone, F. Boulanger & G. Duvert (Dordrecht: Kluwer Academic Publishers), 119
 Falgarone, E., Puget, J.-L., & Péroult, M. 1992, A&A, 257, 715

- Falgarone, E., Lis, D. C., Phillips, T. G., Pouquet, A., Porter, D. H., and Woodward, P. R. 1994, *ApJ*, 436, 728
- Fleck, R. C. 1983, *ApJ*, 272, L45
- Franco, J., Tenorio-Tagle, G., Bodenheimer, P., Rozyczka, M., & Mirabel, I. F. 1988, *ApJ*, 333, 826
- Frisch, U. & Sornette, D. 1997, *Jour. de Phys. I*, 7, 1155
- Genzel, R., Reid, M. J., Moran, J. M., & Downes, D. 1981, *ApJ*, 244, 884
- Frerking, M. A., Langer, W. D., & Wilson, R. W. 1982, *ApJ*, 262, 590
- Frerking, M. A., Keene, J., Blake, G. A. & Phillips, T. G. 1989, *ApJ*, 344, 311
- Genzel, R. & Stutzki, J. 1989, *ARA&A*, 27, 41
- Giannakopoulou, J., Mitchell, G. F., Hasegawa, T. I., Matthews, H. E., & Maillard, J.-P. 1997, *ApJ*, 487, 346
- Goodman, A. A., Benson, P. J., Fuller, G. A., & Myers, P. C. 1993, *ApJ*, 406, 528
- Goodman, A. A., Barranco, J. A., Wilner, D. J., and Heyer, M. H. 1998, *ApJ*, 504, 223
- Graham, J. A. & Heyer, M. 1990, *PASP*, 102, 972
- Graham, J. A. & Rubin, V. C. 1992, *PASP*, 104, 730
- Gregorio, J., Montmerle, T., Casanova, S., & Feigelson, E. D. 1998, *A&A*, 331, 193
- Grenier, I. A., Lebrun, F., Arnaud, M., Dame, T. M., and Thaddeus, P. 1989, *ApJ*, 347, 231
- Gill, A. G. & Henriksen, R. H. 1990, *ApJ*, 365, L27
- Gonatas, C. P., Palmer, P., & Novak, G. 1992, *ApJ*, 398, 118
- Haikala, L. K. & Laureijs, R. J. 1989, *A&A*, 223, 287
- Hobbs, L. M. 1969, *ApJ*, 157, 135
- . 1969, *ApJ*, 157, 165
- . 1974, *ApJ*, 191, 395
- Hobson, M. P. 1992, *MNRAS*, 256, 457
- Houlahan, P. & Scalo, J. M. 1992, *ApJ*, 393, 172
- Huang, S. S. 1950, *ApJ*, 112, 399
- Jayesh & Warhaft, Z. 1991, *Phys. Rev. Lett.*, 67, 3503
- Kailasnath, P., Sreenivasan, K. R., & Stolovitzky, G. 1992, *Phys. Rev. Lett.*, 68, 2766
- Kaplan, S. A. 1954, *Astr. J. U.S.S.R.*, 31, 137
- . 1966, *Interstellar Gas Dynamics*, 2nd edn. (Oxford: Pergamon)
- Keto, E. R. & Lattanzio, J. C. 1989, *ApJ*, 346, 184
- Kida, S. & Murakami, Y. 1989, *Fluid Dyn. Res.*, 4, 347
- Kim, S., Staveley-Smith, L., Dopita, M. A., Freeman, K. C., Sault, R. J., Kesteven, M. J., and McConnell, D. 1998, *ApJ*, 503, 674
- Kitamura, Y., Sunada, K., Hayashi, M., & Hasegawa, T. 1993, *ApJ*, 413, 221
- Kofman, L., Bertschinger, E., Gelb, J. M., Nusser, A., & Dekel, A. 1994, *ApJ*, 420, 44
- Kolmogorov, A. N. 1941, *Dokl. Akad. Nauk SSSR*, 30, 301 (Reprinted in *Proc. R. Soc. Lond. A*, 434, 9 (1991))
- Kornreich, P. & Scalo, J. 1999, *ApJ*, in press (astro-ph/9708131)
- Kulkarni, S. R. & Fich, M. 1985, *ApJ*, 289, 792
- Kramer, C., Stutzki, J., & Winnewisser, G. 1996, 307, 915
- Lada, C. J., Lada, E. A., Clemens, D. P., & Bally, J. 1994, *ApJ*, 429, 694
- Lada, E. A., Bally, J., & Stark, A. A. 1991, *ApJ*, 368, 432
- Lada, E. A., DePoy, D. L., Evans, N. J., & Gatley, I. 1991, *ApJ*, 371, 171
- Lada, E. A., Bally, J., & Stark, A. A. 1992, *ApJ*, 393, L25
- Lamballaise, E., Lesieur, M., & Matais, O. 1997, *Phys. Rev. E*, 56, 6761
- Langer, W. D., Wilson, R. W. & Anderson, C. H. *ApJ*, 408, L45
- Lesieur, M. 1990, *Turbulence in Fluids: Stochastic and Numerical Modeling* (Dordrecht: Kluwer Academic Publishers)
- Lis, D. C., Pety, J., Phillips, T. G., & Falgarone, E. 1996, *ApJ*, 463, 623
- Lis, D. C., Keene, J., Li, Y., Phillips, T. G., and Pety, J. 1998, *ApJ*, in press
- Lis, D. C., Phillips, T. G., Gerin, M., Keene, J., Li, Y., Pety, J., & Falgarone, E. 1998, in *Interstellar Turbulence*, ed. J. Franco & A. Carraminana (Cambridge: Cambridge Univ. Press)
- Little, L. T., Heaton, B. D., & Dent, W. R. F. 1990, *A&A*, 232, 173
- Lockman, F. J., & Gehman, C. S. 1991, *ApJ*, 382, 182
- Lohse, D. & Grossmann, S. 1993, *Physica A*, 194, 519
- Loren, R. B. 1977, *A&A*, 232, 173
- L'vov, V. & Procaccia, I. 1997, in *Turbulence Modeling and Vortex Dynamics*, ed. O. Boratav, A. Eden, & A. Erzan (Berlin: Springer), 1
- Lynds, B. T. 1962 *ApJS*, 7, 1
- Maddalena, R. J., Morris, M., Moscowitz, J., & Thaddeus, P. 1986, *ApJ*, 303, 375
- Mast, J. W. & Goldstein, S. J. 1970, *ApJ*, 159, 319 (MG70)
- Meyers-Rice, B. A. & Lada, C. J. 1991, *ApJ*, 368, 445
- Miesch, M. S. & Bally, J. 1994, *ApJ*, 429, 645 (MB)
- Miesch, M. S. & Scalo, J. M. 1994, *Bull. A.A.S.*, 26, 1458
- . 1995, *ApJ*, 450, L27 (MS)
- Miesch, M. S., Scalo, J. M., & Bally, J. 1999, in preparation (Paper II)
- Miville-Deschenes, M.-A., Joncas, G., & Falgarone, E. 1998, preprint (astro-ph/9804209)
- Monin, A. S. & Yaglom, A. M. 1971, *Statistical Fluid Mechanics: Mechanics of Turbulence* (Cambridge: MIT Press)
- Montalban, J., Bachiller, R., Martinpintado, J., Tafalla, M., GomezGonzalez, J. 1990, *A&A*, 233, 527
- Moriarty-Schieven, G. H. & Snell, R. L. 1988, *ApJ*, 332, 364
- Munch, G. 1957, *ApJ*, 132, 319
- Mundt, R., Ray, T. P., & Buhrke, T. 1988, *ApJ*, 333, L69
- Mundt, R., Ray, T. P., Buhrke, T., Raga, A. C., & Solf, J. 1990, *A&A*, 232, 37
- Myers, P. C. & Khersonsky, V. K. 1995, *ApJ*, 442, 186
- Nakano, M., Sugitani, K., Sato, F. & Ogura, K. 1994, *ApJ*, 423, L147
- Oey, M. S. & Clarke, C. J. 1998, in *Interstellar Turbulence*, ed. J. Franco and A. Carraminana (Cambridge University Press), in press
- Ogura, K. & Sato, F. 1990, *PASJ*, 42, 583
- Olofsson, G. & Koornneef, J. 1985, *A&A*, 146, 337
- Padoan, P., Bally, J., Billawala, Y., Juvela, M., & Nordlund, Å, preprint, available at <http://xxx.lanl.gov/abs/astro-ph/9711008>
- Peebles, P. J. E. 1976, *Ap&SS*, 45, 3
- Pound, M. W., Bania, T. M., & Wilson, R. W. 1990, *ApJ*, 351, 165
- Pound, M. W., & Bally, J. 1991, *ApJ*, 383, 705
- Porter, D. H., Pouquet, A., & Woodward, P. R. 1994, *Phys. Fluids*, 6, 2133
- Pumir, A. 1996, *Phys. Fluids*, 8, 3112
- Racine R. 1968, *AJ*, 73, 233
- Reipurth, B. 1985, *A&A Suppl. Ser.*, 61, 319
- . 1989, *A&A*, 220, 249
- Reipurth, B. & Olberg, M. 1991, *A&A*, 246, 535
- Robinson, L. J. 1984, *S&T*, 67, 4
- Rodriguez, L. F., Moran, J. M., Ho, P. T. P., & Gottlieb, E. W. 1980, *ApJ*, 235, 845
- Scalo, J. M. 1990, in *Physical Processes in Fragmentation and Star Formation*, ed. R. Capuzzo-Dolcetta, C. Chiosi & A. DiFazio (Dordrecht: Kluwer Academic Press), 151
- . 1998, in *The Birth of Galaxies*, XII Blois Conference, in press.
- Scalo, J. M. & Chappell, D. 1998, in preparation
- Scalo, J. M., Chappell, D., & Miesch, M. S. 1999, in preparation (Paper III)
- Seto, N. & Yokoyama, J. 1998, *ApJ*, 492, 421
- She, Z.-S. 1991, *Fluid Dyn. Res.*, 8, 143
- . 1997, in *Turbulence Modeling and Vortex Dynamics*, ed. O. Boratav, A. Eden, & A. Erzan (Berlin: Springer), 28
- She, Z.-S., Chen, S., Doolen, G., Kraichnan, R. H., & Orszag, S. A. 1993, *Phys. Rev. Lett.*, 70, 3251
- Silk, J. 1995, *ApJ*, 438, L41
- Siluk, R. S. & Silk, J. 1974, *ApJ*, 192, 51
- Snell, R. L. 1981, *ApJS*, 45, 121
- Snell, R. L., Loren, R. B., & Plambeck, R. L. 1980, *ApJ*, 239, L17
- Stoeke, J. T., Hartigan, P. M., Strom, S. E., Strom, K. M., Anderson, E. R., Hartmann, L. W., & Kenyon S. J. 1988, *ApJS*, 68, 229
- Stutzki, J., Bensch, F., Heithausen, A., Ossenkopf, V., & Zielinsky, M. 1998, *A&A*, in press
- Tabeling, P. et al. 1996, *Phys. Rev.*, E53, 1613
- Tafalla, M., Bachiller, R., Wright, M. C. H., & Welch, W. J. 1997, *ApJ*, 474, 329
- Tafalla, M. & Myers, P. C. 1997, *ApJ*, 491, 653
- Takakubo, K. 1967, *Bull. Astron. Inst. Netherlands*, 19, 125 (T67)
- Thronson, H. A., Gatley, I., Harvey P. M., Sellgren, K. & Werner, M. W. 1980, *ApJ*, 237, 66
- Torrelles, J. M., Ho, P. T. P., Rodriguez, L. F., Canto, J. 1990, *ApJ*, 349, 529
- Uchida, Y., Kaifu, N., Shibata, K., Hayashi, S. S., Hasegawa, T., & Hamatake, H. 1987, *PASJ*, 39, 907
- Ueda, H., Itoh, M., & Sato, Y. 1993, *ApJ*, 408, 3
- Ungerechts, H. & Thaddeus, P. 1987, *ApJS*, 63, 645
- VanAtta, C. W. & Antonia, R. A. 1980, *Phys. Fluids*, 23, 252
- Verschuur, G. L. 1974, *ApJS*, 27, 65 (V74)
- Vincent, A. & Meneguzzi, M. 1991, *J. Fluid Mech.*, 225, 1
- Vio, R., Fasano, G., Lazzarin, M., & Lessi, O. 1994, *A&A*, 289, 640
- Wiseman, J. J. & Adams, F. C. 1994, *ApJ*, 435, 708
- Wolf, G. A., Lada, C. J., & Bally, J. 1990, *AJ*, 100, 1892
- Wood, D. O., Myers, P. C., & Daugherty, D. A. 1994, *ApJS*, 95, 457
- Wright, J. & Schult, R. L. 1993, *Chaos*, 3, 295
- Xie, T. & Goldsmith, P. F. 1994, *ApJ*, 430, 252
- Xie, T., Goldsmith, P. F., & Patel, N. 1993, *ApJ*, 419, L33
- Yee, E. & Chan, R. 1997, *Phys. Lett. A*, 227, 340
- Yonekura, Y., Dobashi, K., Mizuno, A., Ogawa, H., & Fukuwi, Y. 1997, *ApJS*, 110, 21
- Zimmermann, T. & Stutzki, J. 1993, *Fractals*, 1, 930

This figure "fig01.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig02.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig03.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig04.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig05.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig06.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig07.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig08.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig09.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig10.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig11.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig12.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig13.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig14.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>

This figure "fig15.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810427v2>